

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
24 January 2002 (24.01.2002)

PCT

(10) International Publication Number
WO 02/06223 A1

(51) International Patent Classification⁷: C07D 207/12, 207/14, 207/16, 401/12, 401/14, 403/12, 405/12, 409/12, 417/12, 403/04, 417/14, A61K 31/4025, A61P 9/10, 25/28, 29/00

(21) International Application Number: PCT/US01/21929

(22) International Filing Date: 11 July 2001 (11.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data: 09/615,151 13 July 2000 (13.07.2000) US

(71) Applicant: ABBOTT LABORATORIES [US/US]; Dept. 377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-6050 (US).

(72) Inventors: BENNANI, Youssef, L.; 28572 Windmere Court, Lake Bluff, IL 60044 (US). FAGHIH, Ramin; 36 North Washington Road, Lake Forest, IL 60045 (US). DWIGHT, Wesley, J.; 1943 5th Avenue, San Diego, CA 92101 (US). VASUDEVAN, Anil; 2005 Greystem Circle #308, Gurnee, IL 60031 (US). CONNER, Scott, E.; 12461 Willow Bend, Elizabethtown, IN 47232 (US).

(74) Agents: COLLINS, Daniel, W. et al.; Dept. 377/AP6D-2, 100 Abbott Park Road, Abbott Park, IL 60064-6050 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: 1,3-DISUBSTITUTED AND 1,3,3-TRISUBSTITUTED PYRROLIDINES AS HISTAMINE-3 RECEPTOR LIGANDS AND THEIR THERAPEUTIC APPLICATIONS

WO 02/06223 A1



(57) Abstract: Compounds of formula (I) are useful in treating diseases or conditions prevented by or ameliorated with histamine-3 receptor ligands. Also disclosed are histamine-3 receptor ligand compositions and methods of antagonizing or agonizing histamine-3 receptors.

1,3-DISUBSTITUTED AND 1,3,3-TRISUBSTITUTED PYRROLIDINES AS
HISTAMINE-3 RECEPTOR LIGANDS AND THEIR THERAPEUTIC APPLICATIONS

5 Technical Field

This invention relates to compounds of formula I, which may be useful for treating diseases or conditions caused by or exacerbated by histamine-3 receptor activity, pharmaceutical compositions containing compounds of formula I and methods of treatment using compounds of formula I.

10

Background of the Invention

Histamine is a well-known mediator in hypersensitive reactions (e.g. allergies, hay fever, and asthma) which are commonly treated with antagonists of histamine or "antihistamines." It has also been established that histamine receptors exist in at least two distinct types, referred to as H₁ and H₂ receptors.

15

A third histamine receptor (H₃ receptor) is believed to play a role in neurotransmission in the central nervous system, where the H₃ receptor is thought to be disposed presynaptically on histaminergic nerve endings (Nature, 302, 832-837 (1983)). The existence of the H₃ receptor has been confirmed by the development of selective H₃ receptor agonists and antagonists (Nature, 327, 117-123 (1987)) and has subsequently been shown to regulate the release of other neurotransmitters in both the central nervous system and peripheral organs, particularly the lungs, cardiovascular system and gastrointestinal tract.

20

A number of diseases or conditions may be treated with histamine-3 receptor ligands wherein the H₃ ligand may be an antagonist, agonist or partially agonist, see: (Imamura et al., Circ.Res., (1996) 78, 475-481); (Imamura et. al., Circ.Res., (1996) 78, 863-869); (Lin et al., Brain Res. (1990) 523, 325-330); (Monti et al., Neuropsychopharmacology (1996) 15, 31-35); (Sakai, et al., Life Sci. (1991) 48, 2397-2404); (Mazurkiewicz-Kwilecki and Nsonwah, Can. J. Physiol. Pharmacol. (1989) 67, 75-78); (Panula, P. et al., Neuroscience (1998) 44, 465-481); (Wada et al., Trends in Neuroscience (1991) 14, 415); (Monti et al., Eur. J. Pharmacol. (1991) 205, 283); (Mazurkiewicz-Kwilecki and Nsonwah, Can. J. Physiol. Pharmacol. (1989) 67, 75-78); (Haas et al., Behav. Brain Res. (1995) 66, 41-44); (De Almeida and Izquierdo, Arch. Int. Pharmacodyn. (1986) 283, 193-198); (Kamei et al.,


Psychopharmacology (1990) 102, 312-318); (Kamei and Sakata, Jpn. J. Pharmacol. (1991) 57, 437-482); (Schwartz et al., Psychopharmacology; The fourth Generation of Progress. Bloom and Kupfer (eds). Raven Press, New York, (1995) 397); (Shaywitz et al., Psychopharmacology (1984) 82, 73-77); (Dumery and Blozovski, Exp. Brain Res. (1987) 67, 61-69); (Tedford et al., J. Pharmacol. Exp. Ther. (1995) 275, 598-604); (Tedford et al., Soc. Neurosci. Abstr. (1996) 22, 22); (Yokoyama et al., Eur. J. Pharmacol. (1993) 234, 129); (Yokoyama and Iinuma, CNS Drugs (1996) 5, 321); (Onodera et al., Prog. Neurobiol. (1994) 42, 685); (Leurs and Timmerman, Prog. Drug Res. (1992) 39, 127); (The Histamine H₃ Receptor, Leurs and Timmerman (eds), Elsevier Science, Amsterdam, The Netherlands (1998); (Leurs et al., Trends in Pharm. Sci. (1998) 19, 177-183); (Phillips et al., Annual Reports in Medicinal Chemistry (1998) 33, 31-40); (Matsubara et al., Eur. J. Pharmacol. (1992) 224, 145); (Rouleau et al., J. Pharmacol. Exp. Ther. (1997) 281, 1085); (Adam Szelag, "Role of histamine H₃-receptors in the proliferation of neoplastic cells in vitro", Med. Sci. Monit., 4(5): 747-755, (1998)); (Fitzsimons, C., H. Duran, F. Labombarda, B. Molinari and E. Rivera, "Histamine receptors signalling in epidermal tumor cell lines with H-ras gene alterations", Inflammation Res., 47 (Suppl 1): S50-S51, (1998)); (R. Leurs, R.C. Vollinga and H. Timmerman, "The medicinal chemistry and therapeutic potentials of ligand of the histamine H₃ receptor", Progress in Drug Research 45: 170-165, (1995)); (R. Levi and N.C.E. Smith, "Histamine H₃-receptors: A new frontier in myocardial ischemia", J. Pharm. Exp. Ther., 292: 825-830, (2000)); (Hatta, E., K. Yasuda and R. Levi, "Activation of histamine H₃ receptors inhibits carrier-mediated norepinephrine release in a human model of protracted myocardial ischemia", J. Pharm. Exp. Ther., 283: 494-500, (1997); (H. Yokoyama and K. Iinuma, "Histamine and Seizures: Implications for the treatment of epilepsy", CNS Drugs, 5(5): 321-330, (1995)); (K. Hurukami, H. Yokoyama, K. Onodera, K. Iinuma and T. Watanabe, AQ-0145, "A newly developed histamine H₃ antagonist, decreased seizure susceptibility of electrically induced convulsions in mice", Meth. Find. Exp. Clin. Pharmacol., 17(C): 70-73, (1995); (Delaunois A., Gustin P., Garbarg M., and Ansay M., "Modulation of acetylcholine, capsaicin and substance P effects by histamine H₃ receptors in isolated perfused rabbit lungs", European Journal of Pharmacology 277(2-3):243-50, (1995)); and (Dimitriadou, et al., "Functional relationship between mast cells and C-sensitive nerve fibres evidenced by histamine H₃-receptor modulation in rat lung and spleen", Clinical Science.

87(2):151-63, (1994). Such diseases or conditions include cardiovascular disorders such as acute myocardial infarction; memory processes, dementia and cognition disorders such as Alzheimer's disease and attention-deficit hyperactivity disorder; neurological disorders such as Parkinson's disease, schizophrenia, depression, epilepsy, and seizures or convulsions; 5 cancer such as cutaneous carcinoma, medullary thyroid carcinoma and melanoma; respiratory disorders such as asthma; sleep disorders such as narcolepsy; vestibular dysfunction such as Meniere's disease; gastrointestinal disorders, inflammation, migraine, motion sickness, obesity, pain, and septic shock.

WO 00/06254 describes non-imidazole alkylamines as histamine-3 receptor ligands. 10 EP 0 978 512 A1 describes non-imidazole aryloxy alkylamines as histamine-3 receptor ligands.

Summary of the Invention


In its principle embodiment, the present invention discloses compounds of formula I:



or a pharmaceutically acceptable salt thereof, wherein

Z is selected from the group consisting of a covalent bond and CH₂;

R₁ is selected from the group consisting of OR₂, NR₃R₄ and



20 R₂ is selected from the group consisting of hydrogen, alkoxy carbonyl, alkyl, alkyl carbonyl, aminocarbonyl, sulfono and phosphono;

R₃ and R₄ are independently selected from the group consisting of hydrogen, alkenyl, alkenyl carbonyl, alkenyloxycarbonyl, alkenylsulfonyl, alkoxy carbonyl, alkyl, alkyl carbonyl, 25 alkylsulfonyl, alkynyl, alkynyl carbonyl, alkynyloxycarbonyl, alkynylsulfonyl, aminocarbonyl, aminosulfonyl, arylalkyl, arylalkenyl carbonyl, arylalkenylsulfonyl,

arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylheterocyclecarbonyl, arylheterocylesulfonyl, aryloxyarylcarbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkylcarbonyl, cycloalkylalkylsulfonyl, cycloalkylcarbonyl, cycloalkylsulfonyl, formyl, heterocycle, heterocyclealkyl, 5 heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcarbonyl, heterocyclearylsulfonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcarbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl;

10 R_5 and R_6 are independently selected from the group consisting of hydrogen and alkyl;

R_7 is selected from the group consisting of hydrogen and alkyl; or

R_1 and R_7 together form (=O);

15 R_8 is selected from the group consisting of alkylcarbonyl, aryl, arylcarbonyl, arylcarbonylaryl, arylcarbonylheterocycle, cycloalkylcarbonyl, cycloalkylcarbonylaryl, cycloalkylcarbonylheterocycle, heterocycle, heterocyclecarbonyl, heterocyclecarbonylaryl, heterocyclecarbonylheterocycle; and


R_9 is selected from the group consisting of hydrogen and lower alkyl.

Detailed Description of the Invention

20 All patents, patent applications, and literature references cited in the specification are herein incorporated by reference in their entirety. In the case of inconsistencies, the present disclosure, including definitions, will prevail.

25 It is understood that the foregoing detailed description and accompanying examples are merely illustrative and are not to be taken as limitations upon the scope of the invention, which is defined solely by the appended claims and their equivalents. Various changes and modifications to the disclosed embodiments will be apparent to those skilled in the art. Such changes and modifications, including without limitation those relating to the chemical structures, substituents, derivatives, intermediates, syntheses, formulations and/or methods of use of the invention, may be made without departing from the spirit and scope thereof.

In its principle embodiment, the present invention discloses compounds of formula I:




I,

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof, wherein

5 Z is selected from a covalent bond and CH₂;

R₁ is selected from OR₂, NR₃R₄ and



R₂ is selected from the group consisting of hydrogen, alkoxy carbonyl, alkyl,

alkyl carbonyl, aminocarbonyl, sulfono and phosphono; R₃ and R₄ are independently selected

10 from hydrogen, alkenyl, alkenyl carbonyl, alkenyl oxycarbonyl, alkenyl sulfonyl, alkoxy carbonyl, alkyl, alkyl carbonyl, alkyl sulfonyl, alkynyl, alkynyl carbonyl, alkynyl oxycarbonyl, alkynyl sulfonyl, aminocarbonyl, aminosulfonyl, aryl alkyl, aryl alkenyl carbonyl, aryl alkenyl sulfonyl, aryl alkyl carbonyl, aryl alkyl sulfonyl, aryl aryl carbonyl, aryl aryl sulfonyl, aryl carbonyl, aryl heterocycle carbonyl,

15 aryl heterocycle sulfonyl, aryl oxy aryl carbonyl, aryl oxy aryl sulfonyl, aryl sulfonyl, cyclo alkyl, cyclo alkyl alkyl, cyclo alkyl alkyl carbonyl, cyclo alkyl alkyl sulfonyl, cyclo alkyl carbonyl, cyclo alkyl sulfonyl, formyl, heterocycle, heterocycle alkyl, heterocycle alkyl carbonyl, heterocycle alkyl sulfonyl, heterocycle aryl carbonyl, heterocycle aryl sulfonyl,

20 heterocycle carbonyl, heterocycle heterocycle carbonyl, heterocycle heterocycle sulfonyl, heterocycle oxy alkyl carbonyl, heterocycle oxy aryl carbonyl, heterocycle oxy aryl sulfonyl, heterocycle sulfonyl, and heterocycle thio alkyl carbonyl;

R₅ and R₆ are independently selected from hydrogen and alkyl;

R₇ is selected from hydrogen and alkyl; or

R₁ and R₇ together form (=O);

25 R₈ is selected from the group consisting of alkyl carbonyl, aryl, aryl carbonyl, aryl carbonyl aryl, aryl carbonyl heterocycle, cyclo alkyl carbonyl, cyclo alkyl carbonyl aryl,

cycloalkylcarbonylheterocycle, heterocycle, heterocyclecarbonyl, heterocyclecarbonylaryl, heterocyclecarbonylheterocycle;

R₉ is selected from the group consisting of hydrogen and lower alkyl;

wherein at each occurrence of said aryl, arylalkenylcarbonyl, arylalkenylsulfonyl, 5 arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylcarbonylaryl, arylcarbonylheterocycle, arylheterocyclecarbonyl, arylheterocylesulfonyl, aryloxyarylcarbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkylcarbonylaryl, heterocyclearylcarbonyl, heterocyclearylsulfonyl, heterocyclecarbonylaryl, heterocycleoxyarylcarbonyl, and heterocycleoxyarylsulfonyl, the aryl portion can be 10 optionally substituted with 1, 2, or 3 substituents selected from the group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, aminoalkyl, aminocarbonyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, halogen, haloalkoxy, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and nitro; and

15 wherein at each occurrence of said arylcarbonylheterocycle, arylheterocyclecarbonyl, arylheterocylesulfonyl, cycloalkylcarbonylheterocycle, heterocycle, heterocyclealkyl, heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcarbonyl, heterocyclearylsulfonyl, heterocyclecarbonyl, heterocyclecarbonylaryl, heterocyclecarbonylheterocycle, heterocycleheterocyclecarbonyl, 20 heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcarbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl, the heterocycle portion can be optionally substituted with 1, 2, or 3 substituents selected from the group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, 25 aminoalkyl, aminocarbonyl, aminosulfonyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, halogen, haloalkoxy, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and nitro.

In a preferred embodiment, compounds of the present invention have formula I wherein R₁ is NR₃R₄; R₃ and R₄ are independently selected from hydrogen, alkenyl, alkenylcarbonyl, alkenyloxycarbonyl, alkenylsulfonyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, alkynyl, alkynylcarbonyl, alkynyloxycarbonyl, alkynylsulfonyl, 30 aminocarbonyl, aminosulfonyl, arylalkyl, arylalkenylcarbonyl, arylalkenylsulfonyl,

arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylheterocyclecarbonyl, arylheterocylesulfonyl, aryloxyarylcarbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkylcarbonyl, cycloalkylalkylsulfonyl, cycloalkylcarbonyl, cycloalkylsulfonyl, heterocycle, heterocyclealkyl, 5 heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcarbonyl, heterocyclearylsulfonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcarbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl; R₇ is hydrogen; R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, 10 heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and Z and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula I wherein R₁ is NR₃R₄; R₃ and R₄ are independently selected from hydrogen, alkenyloxycarbonyl, alkoxy carbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, aminocarbonyl, 15 aminosulfonyl, arylalkenylsulfonyl, arylcarbonyl, arylsulfonyl, cycloalkylcarbonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl; R₇ is hydrogen; R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and Z and R₉ are as defined in 20 formula I.


In another preferred embodiment, compounds of the present invention have formula I wherein Z is CH₂; R₁ is NR₃R₄; R₃ and R₄ are independently selected from hydrogen, alkoxy carbonyl and alkyl; R₇ is hydrogen; R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and R₉ is hydrogen. 25

30 In another preferred embodiment, compounds of the present invention have formula I wherein Z is a covalent bond; R₁ is NR₃R₄; R₃ and R₄ are independently selected from

hydrogen, alkoxy carbonyl and alkyl; R₇ is hydrogen; R₈ is selected from the group consisting of acetyl, propionyl, cyclopropyl carbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinyl carbonyl)phenyl, 4-(1-pyrrolidinyl carbonyl)-1-piperazinyl, 4-(1-piperidinyl carbonyl)phenyl, 4-(1-piperidinyl carbonyl)-1-piperazinyl, 4-(4-morpholinyl carbonyl)phenyl, 4-(4-morpholinyl carbonyl)-1-piperazinyl, 4-(1-azetidinyl carbonyl)phenyl, 4-(1-azetidinyl carbonyl)-1-piperazinyl, 4-(1-piperazinyl carbonyl)phenyl, 4-(1-piperazinyl carbonyl)-1-piperazinyl; and R₉ is hydrogen.

In another preferred embodiment, compounds of the present invention have formula II

10



II,

15

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein R₁₀ is selected from alkenyl, alkyl, alkynyl, amino, aryl, arylalkenyl, arylalkyl, arylaryl, arylheterocycle, aryloxyaryl, cycloalkyl, cycloalkylalkyl, heterocycle, heterocyclealkyl, heterocyclearyl, heterocycleheterocycle and heterocycleoxyaryl; and Z, R₃, R₈ and R₉ are as defined in formula I.

20

In another preferred embodiment, compounds of the present invention have formula II wherein R₃ is selected from hydrogen and alkyl; R₈ is selected from alkyl carbonyl, cycloalkyl carbonyl, aryl, heterocycle, heterocycle carbonyl aryl and heterocycle carbonyl heterocycle; R₁₀ is selected from alkyl, amino, aryl, arylalkenyl, heterocycle, heterocycle heterocycle and heterocycle oxyaryl; and Z and R₉ are as defined in formula I.

25

In another preferred embodiment, compounds of the present invention have formula II wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropyl carbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinyl carbonyl)phenyl, 4-(1-pyrrolidinyl carbonyl)-1-piperazinyl, 4-(1-piperidinyl carbonyl)phenyl, 4-(1-piperidinyl carbonyl)-1-piperazinyl, 4-(4-morpholinyl carbonyl)phenyl, 4-(4-morpholinyl carbonyl)-1-piperazinyl, 4-(1-

azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is selected from alkyl and amino.

In another preferred embodiment, compounds of the present invention have formula II
5 wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is aryl wherein said aryl is phenyl optionally substituted with 1 or 2 substituents selected from alkoxy, alkyl, alkylsulfonyl, amino, cyano, haloalkoxy, haloalkyl, halogen.

In another preferred embodiment, compounds of the present invention have formula II
15 wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is arylalkenyl wherein the aryl portion of said arylalkenyl is phenyl.

In another preferred embodiment, compounds of the present invention have formula II
wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl,
25 propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is heterocycle wherein said heterocycle is selected from the group consisting of benzothienyl,

imidazolyl, isoquinolinyl, pyridinyl, pyrrolyl, quinolinyl, thiazolyl, and thienyl, wherein said benzothienyl, imidazolyl, isoquinolinyl, pyridinyl, pyrrolyl, quinolinyl, thiazolyl, and thienyl are each optionally substituted with 1 or 2 substituents selected from alkoxy, alkyl, amino, haloalkyl, and halogen.

5 In another preferred embodiment, compounds of the present invention have formula II wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(1-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is heterocycleheterocycle wherein said heterocycleheterocycle is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)-1H-pyrrol-2-yl.


10 In another preferred embodiment, compounds of the present invention have formula II wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(1-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is heterocycleheterocycle wherein said heterocycleheterocycle is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)-1H-pyrrol-2-yl.

15 In another preferred embodiment, compounds of the present invention have formula II wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(1-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is heterocycleoxyaryl wherein said heterocycleoxyaryl is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)oxy)phenyl).

20 In another preferred embodiment, compounds of the present invention have formula II wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(1-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₀ is heterocycleoxyaryl wherein said heterocycleoxyaryl is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)oxy)phenyl).

25

In another preferred embodiment, compounds of the present invention have formula III



III,

5 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein R₁₁ is selected from hydrogen, alkenyl, alkenyloxy, alkoxy, alkyl, alkynyl, alkynyloxy, amino, aryl, arylalkenyl, arylalkyl, arylaryl, arylheterocycle, aryloxyaryl, cycloalkyl, cycloalkylalkyl, heterocycle, heterocyclealkyl, heterocyclearyl, heterocycleheterocycle, heterocycleoxyalkyl, heterocycleoxyaryl and heterocyclethioalkyl; and Z, R₃, R₈ and R₉ are as defined in formula 10 I.

In another preferred embodiment, compounds of the present invention have formula III wherein R₃ is selected from hydrogen, alkyl, alkylcarbonyl, aminocarbonyl, arylcarbonyl, and heterocyclecarbonyl; R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and R₁₁ is selected 15 from alkenyloxy, alkoxy, alkyl, amino, aryl, cycloalkyl, heterocycle, heterocycleheterocycle and heterocyclethioalkyl; and Z and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula III wherein Z is CH₂; R₃ is selected from hydrogen, alkyl, alkylcarbonyl, and aminocarbonyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₁ 25 is selected from alkenyloxy, alkoxy, alkyl, amino, and cycloalkyl.

In another preferred embodiment, compounds of the present invention have formula III wherein Z is CH₂; R₃ is selected from hydrogen, alkyl, and arylcarbonyl wherein the aryl portion of said arylcarbonyl is phenyl optionally substituted with 1 substituent selected from

cyano and halogen; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₁ is aryl wherein said aryl is phenyl optionally substituted with 1 substituent selected from cyano and halogen.

In another preferred embodiment, compounds of the present invention have formula III wherein Z is CH₂; R₃ is selected from hydrogen, alkyl, and heterocyclecarbonyl wherein the heterocycle portion of said heterocyclecarbonyl is selected from 4-morpholinyl and 1-pyrrolidinyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₁ is heterocycle wherein said heterocycle is selected from furyl, 4-morpholinyl, pyridinyl, and pyrrolidinyl.


In another preferred embodiment, compounds of the present invention have formula III wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₁ is heterocycleheterocycle wherein said heterocycleheterocycle is 2-(3-pyridinyl)-1,3-thiazol-4-yl.

In another preferred embodiment, compounds of the present invention have formula III wherein Z is CH₂; R₃ is selected from hydrogen and alkyl; R₈ is selected from acetyl,

propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₁₁ is heterocyclethioalkyl wherein said heterocyclethioalkyl is [(4-methyl-2-pyrimidinyl)sulfanyl]methyl.

10 In another preferred embodiment, compounds of the present invention have formula

IV



IV,

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein R₃ and R₄ are independently selected from hydrogen, alkenyl, alkyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl; and Z, R₈ and R₉ are as defined in formula I.


15 In another preferred embodiment, compounds of the present invention have formula IV wherein R₃ and R₄ are independently selected from hydrogen and alkyl; R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and Z and R₉ are as defined in formula I.

20 In another preferred embodiment, compounds of the present invention have formula IV wherein Z is CH₂; R₃ and R₄ are independently selected from hydrogen and alkyl; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-

azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and R₉ is hydrogen.

In another preferred embodiment, compounds of the present invention have formula V

5



V,


or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein Z, R₂, R₇, R₈ and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula V
10 wherein R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and Z, R₂, R₇ and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula V
15 wherein Z is CH₂; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₂
20 and R₇ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula VI

25




VI,

or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein Z, R₈ and R₉ are as defined in formula I.

5 In another preferred embodiment, compounds of the present invention have formula VI wherein R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; Z and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula VI wherein Z is CH₂; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and R₉ is hydrogen.

15 15 In another preferred embodiment, compounds of the present invention have formula VII



VII,

20 or a pharmaceutically acceptable salt, ester, amide or prodrug thereof wherein Z, R₅, R₆, R₈ and R₉ are as defined in formula I.

In another preferred embodiment, compounds of the present invention have formula VII wherein R₈ is selected from alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; Z, R₅, R₆ and R₉ are as defined in formula I.

25 25 In another preferred embodiment, compounds of the present invention have formula VII wherein Z is CH₂; R₈ is selected from acetyl, propionyl, cyclopropylcarbonyl, 4-

cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; R₉ is hydrogen; and R₅ and R₆ are as defined in formula I.

Another embodiment of the present invention relates to pharmaceutical compositions comprising a therapeutically effective amount of a compound of formula I-VII or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof in combination with a pharmaceutically acceptable carrier.

Another embodiment of the invention relates to a method of modulating the effects of the histamine-3 receptor by agonism of the histamine-3 receptor in a mammal comprising administering a therapeutically effective amount of a compound of formula I-VII or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof in combination with a pharmaceutically acceptable carrier.

Another embodiment of the invention relates to a method of modulating the effects of the histamine-3 receptor by antagonism of the histamine-3 receptor in a mammal comprising administering a therapeutically effective amount of a compound of formula I-VII or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof in combination with a pharmaceutically acceptable carrier.

Another embodiment of the invention relates to a method of treating acute myocardial infarction, Alzheimer's disease, asthma, attention-deficit hyperactivity disorder, cutaneous carcinoma, depression, epilepsy, inflammation, medullary thyroid carcinoma, melanoma, Meniere's disease, migraine, motion sickness, narcolepsy, obesity, pain, Parkinson's disease, schizophrenia, seizures or septic shock comprising administering a therapeutically effective amount of a compound of formula I-VII or a pharmaceutically acceptable salt, ester, amide, or prodrug thereof in combination with a pharmaceutically acceptable carrier.

Definition of Terms

As used throughout this specification and the appended claims, the following terms have the following meanings:

The term "alkenyl," as used herein, refers to a straight or branched chain hydrocarbon containing from 2 to 10 carbons and containing at least one carbon-carbon double bond formed by the removal of two hydrogens. Representative examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.

The term "alkenylcarbonyl," as used herein, refers to an alkenyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkenylcarbonyl include, but are not limited to, 3-butenoyl, 3-pentenoyl, and 4-pentenoyl.

The term "alkenyloxy," as used herein, refers to an alkenyl group, as defined herein, appended to the parent molecular moiety through an oxy group, as defined herein. Representative examples of alkenyloxy include, but are not limited to, allyloxy, 2-butenyloxy, and 3-butenyloxy.

The term "alkenyloxycarbonyl," as used herein, refers to an alkenyloxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkenyloxycarbonyl include, but are not limited to, allyloxycarbonyl, 2-butenyloxycarbonyl, and 3-butenyloxycarbonyl.

The term "alkenylsulfonyl," as used herein, refers to an alkenyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of alkenylsulfonyl include, but are not limited to, allylsulfonyl, 2-butenylsulfonyl, and 3-butenylsulfonyl.

The term "alkoxy," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through an oxy moiety, as defined herein. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-propoxy, butoxy, tert-butoxy, pentyloxy, and hexyloxy.

The term "alkoxyalkoxy," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through another alkoxy group, as defined herein.

Representative examples of alkoxyalkoxy include, but are not limited to, tert-butoxymethoxy, 2-ethoxyethoxy, 2-methoxyethoxy, and methoxymethoxy.

The term "alkoxyalkyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

5 Representative examples of alkoxyalkyl include, but are not limited to, tert-butoxymethyl, 2-ethoxyethyl, 2-methoxyethyl, and methoxymethyl.

The term "alkoxycarbonyl," as used herein, refers to an alkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkoxycarbonyl include, but are not limited to, methoxycarbonyl, 10 ethoxycarbonyl, and tert-butoxycarbonyl.

The term "alkyl," as used herein, refers to a straight or branched chain hydrocarbon containing from 1 to 10 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, 15 n-heptyl, n-octyl, n-nonyl, and n-decyl.

The term "alkylcarbonyl," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein.

Representative examples of alkylcarbonyl include, but are not limited to, acetyl, 1-oxopropyl, 2,2-dimethyl-1-oxopropyl, 1-oxobutyl, and 1-oxopentyl.

20 The term "alkylcarbonyloxy," as used herein, refers to an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an oxy moiety, as defined herein. Representative examples of alkylcarbonyloxy include, but are not limited to, acyloxy, ethylcarbonyloxy, and tert-butylcarbonyloxy.

The term "alkylsulfinyl," as used herein, refers to an alkyl group, as defined herein, 25 appended to the parent molecular moiety through a sulfinyl group, as defined herein.

Representative examples of alkylsulfinyl include, but are not limited to, methylsulfinyl and ethylsulfinyl.

The term "alkylsulfonyl," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein.

30 Representative examples of alkylsulfonyl include, but are not limited to, ethylsulfonyl, isopropylsulfonyl, and methylsulfonyl.

The term "alkylthio," as used herein, refers to an alkyl group, as defined herein, appended to the parent molecular moiety through a thio moiety, as defined herein. Representative examples of alkylthio include, but are not limited to, methylsulfanyl, ethylsulfanyl, tert-butylsulfanyl, and hexylsulfanyl.

5 The term "alkynyl," as used herein, refers to a straight or branched chain hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited to, acetylenyl, 1-propynyl, 2-propynyl, 3-butynyl, 2-pentynyl, and 1-butynyl.

10 The term "alkynylcarbonyl," as used herein, refers to an alkynyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkynylcarbonyl include, but are not limited to, 3-butynoyl, 3-pentynoyl, and 4-pentynoyl.

15 The term "alkynyloxy," as used herein, refers to an alkynyl group, as defined herein, appended to the parent molecular moiety through an oxy group, as defined herein. Representative examples of alkynyloxy include, but are not limited to, 2-butynyloxy, and 3-butynyloxy.

20 The term "alkynyloxycarbonyl," as used herein, refers to an alkynyloxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkynyloxycarbonyl include, but are not limited to, 2-butynyloxycarbonyl, and 3-butynyloxycarbonyl.

25 The term "alkynylsulfonyl," as used herein, refers to an alkynyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of alkynylsulfonyl include, but are not limited to, 2-butynylsulfonyl, and 3-butynylsulfonyl.

30 The term "amino," as used herein, refers to a -NR_AR_B group wherein R_A and R_B are independently selected from hydrogen, alkyl, alkylcarbonyl, and benzyl. Representative examples of amino include but are not limited to acetylamino, amino, benzylamino, dimethylamino, and methylamino.

The term "aminoalkyl," as used herein, refers to an amino group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

Representative examples of aminoalkyl include, but are not limited, (amino)methyl, (dimethylamino)methyl, 2-(benzylamino)ethyl, and (ethylamino)methyl.

The term "aminocarbonyl," as used herein, refers to an amino group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein.

5 Representative examples of aminocarbonyl include, but are not limited, aminocarbonyl, dimethylaminocarbonyl, benzylaminocarbonyl, and ethylaminocarbonyl.

The term "aminosulfonyl," as used herein, refers to an amino group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein.

Representative examples of aminosulfonyl include, but are not limited, aminosulfonyl, 10 dimethylaminosulfonyl, benzylaminosulfonyl, and ethylaminosulfonyl.

The term "aryl," as used herein, refers to a monocyclic-ring system, or a bicyclic- or a tricyclic- fused ring system wherein one or more of the fused rings are aromatic.

Representative examples of aryl include, but are not limited to, anthracenyl, azulenyl, fluorenyl, indanyl, indenyl, naphthyl, phenyl, and tetrahydronaphthyl.

15 The aryl groups of this invention can be substituted with 1, 2, or 3 substituents independently selected from alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, aminoalkyl, aminocarbonyl, aminosulfonyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, halogen, haloalkoxy, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and 20 nitro.

The term "arylalkenyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an alkenyl group, as defined herein.

Representative examples of arylalkenyl include, but are not limited to, 3-phenyl-1-propenyl, and 2-(2-naphthyl)ethenyl.

25 The term "arylalkenylcarbonyl," as used herein, refers to an arylalkenyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylalkenylcarbonyl include, but are not limited to, 4-phenyl-3-butenoyl, and 3-phenyl-2-propenoyl.

30 The term "arylalkenylsulfonyl," as used herein, refers to an arylalkenyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined

herein. Representative examples of arylalkenylsulfonyl include, but are not limited to, 2-phenylethenylsulfonyl, and 4-phenyl-3-butenylsulfonyl.

The term "arylalkyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

5 Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 3-phenylpropyl, and 2-naphth-2-ylethyl.

The term "arylalkylcarbonyl," as used herein, refers to an arylalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylalkylcarbonyl include, but are not limited to, phenylacetyl, 4-phenylbutanoyl, and 3-phenylpropanoyl.

10 The term "arylalkylsulfonyl," as used herein, refers to an arylalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylalkylsulfonyl include, but are not limited to, (2-phenylethyl)sulfonyl, and (3-phenylpropyl)sulfonyl.

15 The term "arylaryl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through another aryl group, as defined herein. Representative examples of arylaryl include, but are not limited to, (1,1'-biphenyl), and (2'-chloro(1,1'-biphenyl)-3-yl).

20 The term "arylarylcarbonyl," as used herein, refers to an arylaryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylarylcarbonyl include, but are not limited to, (1,1'-biphenyl)carbonyl, and (2'-chloro(1,1'-biphenyl)-3-yl)carbonyl.

25 The term "arylaryl sulfonyl," as used herein, refers to an arylaryl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylaryl sulfonyl include, but are not limited to, (1,1'-biphenyl)sulfonyl, and (2'-chloro(1,1'-biphenyl)-3-yl)sulfonyl.

30 The term "arylcarbonyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylcarbonyl include, but are not limited to, benzoyl, 4-cyanobenzoyl, and naphthoyl.

The term "arylcarbonylaryl," as used herein, refers to an arylcarbonyl group, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of arylcarbonylaryl include, but are not limited to, 4-(benzoyl)phenyl and 4-(benzoyl)naphthyl.

5 The term "arylcarbonylheterocycle," as used herein, refers to an arylcarbonyl group, as defined herein, appended to the parent molecular moiety through a heterocycle group, as defined herein. Representative examples of arylcarbonylheterocycle include, but are not limited to, 4-benzoyl-1-piperazinyl and 1-benzoyl-4-piperidinyl.

10 The term "arylheterocycle," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a heterocycle group, as defined herein. Representative examples of arylheterocycle include, but are not limited to, 5-phenylpyridin-2-yl and 5-(3-chlororphenyl)pyridin-2-yl.

15 The term "arylheterocyclecarbonyl," as used herein, refers to an arylheterocycle group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of arylheterocyclecarbonyl include, but are not limited to, 5-phenylpyridin-2-ylcarbonyl and 5-(3-chlororphenyl)pyridin-2-ylcarbonyl.

20 The term "arylheterocyclesulfonyl," as used herein, refers to an arylheterocycle group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylheterocyclesulfonyl include, but are not limited to, 5-phenylpyridin-2-ylsulfonyl and 5-(3-chlororphenyl)pyridin-2-ylsulfonyl.

The term "aryloxy," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through an oxy moiety, as defined herein. Representative examples of aryloxy include, but are not limited to, phenoxy, naphthoxy, 3-bromophenoxy, 4-chlorophenoxy, 4-methylphenoxy, and 3,5-dimethoxyphenoxy.

25 The term "aryloxyaryl," as used herein, refers to an aryloxy group, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of aryloxyaryl include, but are not limited to, 3-(3-methylphenoxy)phenyl, and 3-(3-bromophenoxy)phenyl.

30 The term "aryloxyarylcarbonyl," as used herein, refers to an aryloxyaryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined

herein. Representative examples of aryloxyarylcarbonyl include, but are not limited to, 3-(3-methylphenoxy)benzoyl, and 3-(3-bromophenoxy)benzoyl.

5 The term "aryloxyarylsulfonyl," as used herein, refers to an aryloxyaryl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of aryloxyarylsulfonyl include, but are not limited to, 3-(3-methylphenoxy)phenylsulfonyl, and 3-(3-bromophenoxy)phenylsulfonyl.

10 The term "arylsulfonyl," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of arylsulfonyl include, but are not limited to, phenylsulfonyl, (4-acetylaminophenyl)sulfonyl, (4-chlorophenyl)sulfonyl, (4-cyanophenyl)sulfonyl, (4-methoxyphenyl)sulfonyl, (4-methylphenyl)sulfonyl, and (4-(tert-butyl)phenyl)sulfonyl.

15 The term "arylthio," as used herein, refers to an aryl group, as defined herein, appended to the parent molecular moiety through a thio moiety, as defined herein. Representative examples of arylthio include, but are not limited to, phenylsulfanyl, naphth-2-ylsulfanyl, and 5-phenylhexylsulfanyl.

The term "carbonyl," as used herein, refers to a -C(O)- group.

The term "carboxy," as used herein, refers to a -CO₂H group.

20 The term "carboxyalkyl," as used herein, refers to a carboxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of carboxyalkyl include, but are not limited to, carboxymethyl, 2-carboxyethyl, and 3-carboxypropyl.

The term "cyano," as used herein, refers to a -CN group.

25 The term "cyanoalkyl," as used herein, refers to a cyano group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of cyanoalkyl include, but are not limited to, cyanomethyl, 2-cyanoethyl, and 3-cyanopropyl.

The term "cycloalkyl," as used herein, refers to a saturated cyclic hydrocarbon group containing from 3 to 8 carbons. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl.

30 The term "cycloalkylalkyl," as used herein, refers to cycloalkyl group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

Representative examples of cycloalkylalkyl include, but are not limited to, cyclopropylmethyl, 2-cyclobutylethyl, cyclopentylmethyl, cyclohexylmethyl, and 4-cycloheptylbutyl.

The term "cycloalkylalkylcarbonyl," as used herein, refers to cycloalkylalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of cycloalkylalkylcarbonyl include, but are not limited to, cyclopropylmethylcarbonyl, 2-cyclobutylethylcarbonyl, cyclopentylmethylcarbonyl, cyclohexylmethylcarbonyl, and 4-cycloheptylbutylcarbonyl.

The term "cycloalkylalkylsulfonyl," as used herein, refers to cycloalkylalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of cycloalkylalkylsulfonyl include, but are not limited to, cyclopropylmethylsulfonyl, 2-cyclobutylethylsulfonyl, cyclopentylmethylsulfonyl, cyclohexylmethylsulfonyl, and 4-cycloheptylbutylsulfonyl.

The term "cycloalkylcarbonyl," as used herein, refers to a cycloalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of cycloalkylcarbonyl include, but are not limited to, cyclopropylcarbonyl, cyclopentylcarbonyl, and cyclohexylcarbonyl.

The term "cycloalkylcarbonylaryl," as used herein, refers to a cycloalkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of cycloalkylcarbonylaryl include, but are not limited to, 4-(cyclopropylcarbonyl)phenyl, 4-(cyclopentylcarbonyl)phenyl, and 4-(cyclohexylcarbonyl)phenyl.

The term "cycloalkylcarbonylheterocycle," as used herein, refers to a cycloalkylcarbonyl group, as defined herein, appended to the parent molecular moiety through a heterocycle group, as defined herein. Representative examples of cycloalkylcarbonylheterocycle include, but are not limited to, 4-(cyclopropylcarbonyl)-1-piperazinyl, 4-(cyclopentylcarbonyl)-1-piperazinyl, and 4-(cyclohexylcarbonyl)-1-piperazinyl.

The term "cycloalkylsulfonyl," as used herein, refers to cycloalkyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein.

Representative examples of cycloalkylsulfonyl include, but are not limited to, cyclopropylsulfonyl, cyclopentylsulfonyl, and cyclohexylsulfonyl.

The term "formyl," as used herein, refers to a -C(O)H group.

The term "halo" or "halogen," as used herein, refers to -Cl, -Br, -I or -F.

5 The term "haloalkoxy," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkoxy group, as defined herein. Representative examples of haloalkoxy include, but are not limited to, chloromethoxy, 2-fluoroethoxy, trifluoromethoxy, and pentafluoroethoxy.

10 The term "haloalkyl," as used herein, refers to at least one halogen, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein.

Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, trifluoromethyl, pentafluoroethyl, and 2-chloro-3-fluoropentyl.

15 The term "heterocycle" or "heterocyclic," as used herein, refers to a monocyclic, bicyclic, or tricyclic ring system. Monocyclic ring systems are exemplified by any 3- or 4-membered ring containing a heteroatom independently selected from oxygen, nitrogen and sulfur; or a 5-, 6- or 7-membered ring containing one, two or three heteroatoms wherein the heteroatoms are independently selected from nitrogen, oxygen and sulfur. The 5-membered ring has from 0-2 double bonds and the 6- and 7-membered ring have from 0-3 double bonds.

20 Representative examples of monocyclic ring systems include, but are not limited to, azetidinyl, azepinyl, aziridinyl, diazepinyl, 1,3-dioxolanyl, dioxanyl, dithianyl, furyl, imidazolyl, imidazolinyl, imidazolidinyl, isothiazolyl, isothiazolinyl, isothiazolidinyl, isoxazolyl, isoxazolinyl, isoxazolidinyl, morpholinyl, oxadiazolyl, oxadiazolinyl, oxadiazolidinyl, oxazolyl, oxazolinyl, oxazolidinyl, piperazinyl, piperidinyl, pyranyl, pyrazinyl, pyrazolyl, pyrazolinyl, pyrazolidinyl, pyridyl, pyrimidinyl, pyridazinyl, pyrrolyl, 25 pyrrolinyl, pyrrolidinyl, tetrahydrofuranlyl, tetrahydrothienyl, tetrazinyl, tetrazolyl, thiadiazolyl, thiadiazolinyl, thiadiazolidinyl, thiazolyl, thiazolinyl, thiazolidinyl, thienyl, thiomorpholinyl, 1,1-dioxidothiomorpholinyl (thiomorpholine sulfone), thiopyranyl, triazinyl, triazolyl, and trithianyl. Bicyclic ring systems are exemplified by any of the above

30 monocyclic ring systems fused to an aryl group as defined herein, a cycloalkyl group as defined herein, or another monocyclic ring system. Representative examples of bicyclic ring systems include but are not limited to, for example, benzimidazolyl, benzothiazolyl,

benzothienyl, benzoxazolyl, benzofuranyl, benzopyranyl, benzothiopyranyl, benzodioxinyl, 1,3-benzodioxolyl, cinnolinyl, indazolyl, indolyl, indolinyl, indolizinyl, naphthyridinyl, isobenzofuranyl, isobenzothienyl, isoindolyl, isoindolinyl, isoquinolinyl, phthalazinyl, pyranopyridyl, quinolinyl, quinolizinyl, quinoxalanyl, quinazolinyl, tetrahydroisoquinolinyl, tetrahydroquinolinyl, and thiopyranopyridyl.

5 The heterocycles of this invention can be substituted with 1, 2, or 3 substituents independently selected from alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, aminoalkyl, aminocarbonyl, aminosulfonyl, carboxy, carboxyalkyl, cyano, 10 cyanoalkyl, formyl, halogen, haloalkoxy, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and nitro.

The term "heterocyclealkyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of heterocyclealkyl include, but are not limited to, pyridin-3-ylmethyl and 2-pyrimidin-2-ylpropyl.

15 The term "heterocyclealkylcarbonyl," as used herein, refers to a heterocyclealkyl, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclealkylcarbonyl include, but are not limited to, (pyridin-3-ylmethyl)carbonyl and (2-(pyrimidin-2-yl)propyl)carbonyl.

20 The term "heterocyclealkylsulfonyl," as used herein, refers to a heterocyclealkyl, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocyclealkylsulfonyl include, but are not limited to, (pyridin-3-ylmethyl)sulfonyl and (2-(pyrimidin-2-yl)propyl)sulfonyl.

25 The term "heterocyclearyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of heterocyclearyl include, but are not limited to, 4-(pyridin-3-yl)phenyl and 4-(pyrimidin-2-yl)phenyl.

30 The term "heterocyclearylcarbonyl," as used herein, refers to a heterocyclearyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclearylcarbonyl include, but are not limited to, 4-(pyridin-3-yl)benzoyl and 4-(pyrimidin-2-yl)benzoyl.

The term "heterocyclearylsulfonyl," as used herein, refers to a heterocyclearyl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocyclearylsulfonyl include, but are not limited to, (4-(pyridin-3-yl)phenyl)sulfonyl and (4-(pyrimidin-2-yl)phenyl)sulfonyl.

5 The term "heterocyclecarbonyl," as used herein, refers to a heterocycle, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclecarbonyl include, but are not limited to, 2-furoyl, morpholin-1-ylcarbonyl, pyridin-3-ylcarbonyl, pyrrolidin-1-ylcarbonyl, and quinolin-3-ylcarbonyl.

10 The term "heterocyclecarbonylaryl," as used herein, refers to a heterocyclecarbonyl, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of heterocyclecarbonylaryl include, but are not limited to, 4-(2-furoyl)phenyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)phenyl and 4-(3-pyridinylcarbonyl)phenyl.

15 The term "heterocyclecarbonylheterocycle," as used herein, refers to a heterocyclecarbonyl, as defined herein, appended to the parent molecular moiety through a heterocycle group, as defined herein. Representative examples of heterocyclecarbonylheterocycle include, but are not limited to, 4-(2-furoyl)-1-piperazinyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl and 4-(3-pyridinylcarbonyl)-1-piperazinyl.

20 The term "heterocycleheterocycle," as used herein, refers to a heterocycle group, as defined herein, appended to the parent molecular moiety through another heterocycle group, as defined herein. Representative examples of heterocycleheterocycle include, but are not limited to, 2-(pyridin-3-yl)thiazo-4-yl and 2-(pyrimidin-2-yl)thiazo-4-yl.

25 The term "heterocycleheterocyclecarbonyl," as used herein, refers to a heterocycleheterocycle group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocycleheterocyclecarbonyl include, but are not limited to, (2-(pyridin-3-yl)thiazo-4-yl)carbonyl and (2-(pyrimidin-2-yl)thiazo-4-yl)carbonyl.

The term "heterocycleheterocyclesulfonyl," as used herein, refers to a heterocycleheterocycle group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocycleheterocyclesulfonyl include, but are not limited to, (2-(pyridin-3-yl)thiazo-4-yl)sulfonyl and (2-(pyrimidin-2-yl)thiazo-4-yl)sulfonyl.

The term "heterocycleoxy," as used herein, refers to a heterocycle group, as defined herein, appended to the parent molecular moiety through an oxy moiety, as defined herein. Representative examples of heterocycleoxy include, but are not limited to, pyrid-3-yloxy and quinolin-3-yloxy.

The term "heterocycleoxyalkyl," as used herein, refers to a heterocycleoxy group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of heterocycleoxyalkyl include, but are not limited to, pyrid-3-yloxymethyl and 2-quinolin-3-yloxyethyl.

The term "heterocycleoxyalkylcarbonyl," as used herein, refers to a heterocycleoxyalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocycleoxyalkylcarbonyl include, but are not limited to, (pyridin-3-yloxymethyl)carbonyl and (2-(quinolin-3-yloxy)ethyl)carbonyl.

The term "heterocycleoxyaryl," as used herein, refers to a heterocycleoxy group, as defined herein, appended to the parent molecular moiety through an aryl group, as defined herein. Representative examples of heterocycleoxyaryl include, but are not limited to, 4-(pyridin-3-yloxy)phenyl and 4-(quinolin-3-yloxy)phenyl.

The term "heterocycleoxyarylcarbonyl," as used herein, refers to a heterocycleoxyaryl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocycleoxyarylcarbonyl include, but are not limited to, 4-(pyridin-3-yloxy)benzoyl and 4-(quinolin-3-yloxy)benzoyl.

The term "heterocycleoxyarylsulfonyl," as used herein, refers to a heterocycleoxyaryl group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocycleoxyarylsulfonyl include, but are not limited to, (4-(pyridin-3-yloxy)phenyl)sulfonyl and (4-(quinolin-3-yloxy)phenyl)sulfonyl.

The term "heterocyclesulfonyl," as used herein, refers to a heterocycle group, as defined herein, appended to the parent molecular moiety through a sulfonyl group, as defined herein. Representative examples of heterocyclesulfonyl include, but are not limited to, (pyridin-3-yl)sulfonyl and (quinolin-8-yl)sulfonyl.

5 The term "heterocyclethio," as used herein, refers to a heterocycle group, as defined herein, appended to the parent molecular moiety through a thio moiety, as defined herein. Representative examples of heterocyclethio include, but are not limited to, pyrid-3-ylsulfanyl and quinolin-3-ylsulfanyl.

10 The term "heterocyclethioalkyl," as used herein, refers to a heterocyclethio group, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of heterocyclethioalkyl include, but are not limited to, pyrid-3-ylsulfanylmethyl, (4-methylpyrimidin-2-yl)sulfanylmethyl, and 2-(quinolin-3-ylsulfanyl)ethyl.

15 The term "heterocyclethioalkylcarbonyl," as used herein, refers to a heterocyclethioalkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of heterocyclethioalkylcarbonyl include, but are not limited to, (pyrid-3-ylsulfanyl)acetyl, ((4-methylpyrimidin-2-yl)sulfanyl)acetyl, and (quinolin-3-ylsulfanyl)acetyl.

20 The term "hydroxy," as used herein, refers to an -OH group.

25 The term "hydroxyalkyl," as used herein, refers to one or two hydroxy groups, as defined herein, appended to the parent molecular moiety through an alkyl group, as defined herein. Representative examples of hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 2,3-dihydroxypropyl and 2-ethyl-4-hydroxyheptyl.

30 The term "lower alkyl," as used herein, is a subset of alkyl as defined herein and refers to a straight or branched chain hydrocarbon group containing from 1 to 4 carbon atoms. Examples of lower alkyl are methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and tert-butyl.

The term "mercapto," as used herein, refers to a -SH group.

35 The term "nitro," as used herein, refers to a -NO₂ group.

The term "oxo," as used herein, refers to a =O moiety.

The term "oxy," as used herein, refers to a -O- moiety.

The term "phosphono," as used herein, refers to a -P(O)(OR_D)₂ group wherein R_D is selected from hydrogen and alkyl, as defined herein. Representative examples of phosphono include, but are not limited to, dimethoxyphosphoryl and diethoxyphosphoryl.

5 The term "sulfinyl," as used herein, refers to a -S(O)- group.

The term "sulfono," as used herein, refers to a -S(O)₂(OR_E) group wherein R_E is selected from alkyl, aryl, and arylalkyl, as defined herein. Representative examples of sulfono include, but are not limited to, methoxysulfonyl, ethoxysulfonyl, (benzyloxy)sulfonyl and phenoxy sulfonyl.

10 The term "sulfonyl," as used herein, refers to a -SO₂- group.

The term "thio," as used herein, refers to a -S- moiety.

Preferred compounds of formula I include, but are not limited to:

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}cyclopropanecarboxamide;

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-[(4-methyl-2-

15 pyrimidinyl)sulfanyl]acetamide;

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}nicotinamide;

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide;

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-cyanobenzenesulfonamide;

20 N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-propanesulfonamide;

1-(4-{3-[(3R)-3-aminopyrrolidinyl]propoxy}phenyl)-1-propanone;

(5R)-3-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-methyl-2,4-imidazolidinedione;

(5S)-3-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-methyl-2,4-

25 imidazolidinedione;

(5R)-3-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-isopropyl-2,4-imidazolidinedione;

4-cyano-N-((3R)-1-{3-[4-

(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

30 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-propanesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide;

N-((3R)-1-{3-[4-acetylphenoxy]propyl}pyrrolidinyl)-4-cyanobenzamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)benzenesulfonamide;

5 4-bromo-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

10 4-chloro-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-(4-{[(3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl]amino}sulfonyl)phenyl)acetamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-methoxybenzenesulfonamide;

15 4-tert-butyl-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

20 N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-2,5-dimethoxybenzenesulfonamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-2-methylbenzenesulfonamide;

25 3-chloro-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-ethylbenzenesulfonamide;

30 N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide;

2-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

5 3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

3,5-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

10 3-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide;

15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-thiophenesulfonamide;

5-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-methyl-1-benzothiophene-2-sulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide;

20 N-(5-{[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl]amino}sulfonyl)-4-methyl-1,3-thiazol-2-yl)acetamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-8-quinolinesulfonamide;

25 4-butoxy-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide;

30 3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-phenylethenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide;

5 2-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-methylbenzenesulfonamide;

10 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide;

15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2,4-difluorobenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-7-isoquinolinesulfonamide;

20 N-(2-chloro-4-{{((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)amino}sulfonyl}phenyl)acetamide;

3,4-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

25 4-bromo-N-((3R)-1-{3-[4-

(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

4-chloro-N-((3R)-1-{3-[4-

30 (cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methoxybenzenesulfonamide;

4-tert-butyl-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

5 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-(trifluoromethyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2,5-dimethoxybenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-methylbenzenesulfonamide;

10 3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

15 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-ethylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide;

20 2-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

25 3-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide;

30 N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-2,4-difluorobenzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-isoquinolinesulfonamide;
N-{4-[{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}amino]sulfonyl]-2-chlorophenyl}acetamide;
N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-3,4-
5 dichlorobenzenesulfonamide;
N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1H-pyrrole-2-sulfonamide;
N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-4-{{3-chloro-5-(trifluoromethyl)-2-pyridinyl}oxy}benzenesulfonamide;
10 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-thiophenesulfonamide;
5-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-
15 methyl-1-benzothiophene-2-sulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-
quinolinesulfonamide;
20 4-butoxy-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide;
3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-
25 methylbenzenesulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-phenylethenesulfonamide;
N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide;
30 2-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-methylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide;

5 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide;

10 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-isoquinolinesulfonamide;

N-(2-chloro-4-{[(3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl]amino}sulfonyl}phenyl)acetamide;
3,4-dichloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

15 tert-butyl 1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}-3-pyrrolidinylcarbamate;
4'-{3-[(3R)-3-aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
tert-butyl (3S)-1-{2-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]ethyl}pyrrolidinylcarbamate;
4-methoxy-N-((3R)-1-{3-[4-(2-

pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

20 4-isopropyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

4-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

25 3-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

1-methyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)-1H-imidazole-4-sulfonamide;

3,4-dimethoxy-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

30 N-(2-chloro-4-{[(3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl]amino}sulfonyl}phenyl)acetamide;

4'-{3-[(3R)-3-(dimethylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
tert-butyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;
4'-{3-[(3R)-3-(methylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
5 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-
methylacetamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,3,3-
trimethylbutanamide;
methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-
10 yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;
tert-pentyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N',N'-
trimethylurea;
15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-1-
pyrrolidinecarboxamide;
N'-tert-butyl-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-
N-methylurea;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-
20 morpholinecarboxamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-
methylbenzamide;
4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-
methylbenzamide;
25 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-
methylnicotinamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-
furamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-(3-
30 pyridinyl)-1,3-thiazole-4-carboxamide;

N-[(3R)-1-[3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl]pyrrolidinyl]-N,N',N'-trimethyl-sulfamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-fluoro-N-methylbenzenesulfonamide;

5 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-isopropyl-N-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-10 (methylsulfonyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-(4-fluorobenzoyl)benzamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)acetamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,3-15 dimethylbutanamide;

allyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate;

methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate;

tert-pentyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate;

20 N'-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N-dimethylurea;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-1-pyrrolidinecarboxamide;

N-(tert-butyl)-N'-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)urea;

25 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-morpholinecarboxamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluorobenzamide;

30 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)nicotinamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-furamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide;
5 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-propanesulfonamide;
N'-(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N-dimethylsulfamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-10 (methylsulfonyl)benzenesulfonamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(3,3-dimethylbutanoyl)-3,3-dimethylbutanamide;
N'-(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N'-(dimethylaminocarbonyl)-N,N-dimethylurea;
15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(1-pyrrolidinylcarbonyl)-1-pyrrolidinecarboxamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(4-morpholinylcarbonyl)-4-morpholinecarboxamide;
cyclopropyl{4-[3-(3-hydroxy-1-pyrrolidinyl)propoxy]phenyl}methanone;
20 cyclopropyl(4-{3-[(3R)-3-hydroxypyrrrolidinyl]propoxy}phenyl)methanone;
4'-{3-[(3R)-3-hydroxypyrrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
4'-[3-(3-oxo-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile;
4'-{3-[(3S)-3-hydroxypyrrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
4'-[3-(3-hydroxy-3-methyl-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile;
25 4'-[3-(3-hydroxy-3-isopropyl-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile;
4'-{3-[(3R)-3-hydroxy-3-methylpyrrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;
N,N-dimethyl-N-[(3R)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl]amine;
N,N-dimethyl-N-[(3S)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-30 yl]oxy}propyl)pyrrolidinyl]amine;

(3R)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)-3-pyrrolidinol; 5 N,N-dimethyl-N-[(3R)-1-(3-{4-[4-(1-pyrrolidinylcarbonyl)-1-piperazinyl]phenoxy}propyl)pyrrolidinyl]amine; N,N-dimethyl-N-[(3S)-1-(3-{4-[4-(1-pyrrolidinylcarbonyl)-1-piperazinyl]phenoxy}propyl)pyrrolidinyl]amine and pharmaceutically acceptable salts, esters, amides, or prodrugs thereof.

Abbreviations

10 Abbreviations which have been used in the descriptions of the schemes and the examples that follow are: AcOH for acetic acid; BF_3OEt_2 for boron trifluoride diethyl ether complex; Boc for tert-butoxycarbonyl; $(\text{Boc})_2\text{O}$ for di-tert-butyl dicarbonate; n-BuLi for n-butyllithium; CDI for 1,1'-carbonyldiimidazole; DCC for 1,3-dicyclohexylcarbodiimide; DMAP for 4-dimethylaminopyridine; DMF for N,N-dimethylformamide; DMSO for 15 dimethyl sulfoxide; EDCI or EDC for 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride; EtOAc for ethyl acetate; EtOH for ethanol; HOBT for 1-hydroxybenzotriazole hydrate; LAH for lithium aluminum hydride; LDA for lithium diisopropylamide; MeOH for methanol; pyr for pyridine; TFA for trifluoroacetic acid; and THF for tetrahydrofuran.

20


Preparation of Compounds of The Invention

The compounds and processes of the present invention will be better understood in connection with the following synthetic schemes and methods which illustrate a means by which the compounds of the invention can be prepared.

25

The compounds of this invention can be prepared by a variety of synthetic routes. Representative procedures are shown in Schemes 1-7.


Scheme 1




Sulfonamides of general formula (6), wherein R_3 and R_8 are as defined in formula I, and R_{10} is selected from alkenyl, alkyl, alkynyl, amino, aryl, arylalkenyl, arylalkyl, arylaryl, arylheterocyle, aryloxyaryl, cycloalkyl, cycloalkylalkyl, heterocyclealkyl, heterocyclearyl, heterocycleheterocycle, heterocycleoxyaryl and heterocycle, may be prepared as described in Scheme 1. Phenols of general formula (1) may be treated with 1-bromo-3-chloropropane (or 1-bromo-2-chloroethane to provide the ethyl analogues) and a base such as potassium carbonate in a solvent such as 2-butanone with heat to provide chlorides of general formula (2). Chlorides of general formula (2) may be treated with tert-butyl pyrrolidinylcarbamate (or tert-butyl (3R)-pyrrolidinylcarbamate or tert-butyl (3S)-pyrrolidinylcarbamate), potassium iodide, a base such as potassium carbonate in a solvent such as 2-butanone with heat to provide N-boc aminopyrrolidines which may be deprotected with acid such as 4N HCl in 1,4-dioxane or trifluoroacetic acid in CH_2Cl_2 to provide aminopyrrolidines of general formula (3). Aminopyrrolidines of general formula (3) may be treated with sulfonyl chlorides of general formula (4), a base such as triethylamine, diisopropylamine or a polymer supported base such as tris(2-aminoethyl)amine-polystyrene resin and catalytic DMAP in a solvent such

as methylene chloride or DMF to provide sulfonamides of general formula (5). Sulfonamides of general formula (5) may be treated with a base such as sodium hydride and an electrophile such as, for example, iodomethane, allyl bromide or propargyl bromide to provide sulfonamides of general formula (6).

5


Scheme 2

Amides of general formula (9), wherein R₃ and R₈ are as defined in formula I and R₁₁ is selected from hydrogen, alkenyl, alkenyloxy, alkoxy, alkyl, alkynyl, alkynyloxy, amino, arylalkenyl, arylalkyl, arylaryl, aryl, arylheterocycle, aryloxyaryl, cycloalkylalkyl, cycloalkyl, heterocyclealkyl, heterocyclearyl, heterocycle, heterocycleheterocycle, heterocycloxyalkyl, heterocycloxyaryl, and heterocylethioalkyl, may be prepared as described in Scheme 2. Aminopyrrolidines of general formula (3), from Scheme 1, may be treated with acids of general formula (7), a coupling reagent such as DCC, EDCI, or a polymer supported coupling reagent (N-cyclohexylcarbodiimide, N'-methyl polystyrene resin), catalytic DMAP, a base such as triethylamine, diisopropylamine or a polymer supported base (tris(2-aminoethyl)amine-polystyrene resin) and optionally HOBT to provide amides of general formula (8). Alternatively, aminopyrrolidines of general formula (3) may be treated with acid chlorides and a base to provide amides of general formula (8). Amides of general formula (8) may be treated with a base such as sodium hydride and an electrophile such as, for example, iodomethane, allyl bromide, propargyl bromide or an acid chloride to provide amides of general formula (9).

Scheme 3


Amines of general formula (10) and (11), wherein R₈ is as defined in formula I and R₃ and R₄ are independently selected from alkenyl, alkyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl, may be prepared as described in Scheme 3. Aminopyrrolidines of general formula (3), from Scheme 1, may be treated with electrophiles such as, for example, iodomethane, allyl bromide, propargyl bromide, benzylbromide, bromocyclohexane or 3,6-dichloropyridazine and a base such as triethylamine or diisopropylamine in a solvent such as DMF or THF to provide monosubstituted amines of 10 general formula (10). Monosubstituted amines of general formula (10) may be retreated with an electrophile and a base to provide disubstituted amines of general formula (11).

Scheme 4




2,5-Dioxo-1-imidazolidines of general formula (15) and (16), wherein R_5 , R_6 , and R_8 are as defined in formula I, may be prepared as described in Scheme 4. Aminopyrrolidines of general formula (3), from Scheme 1, may be treated with N-protected (L) α -amino acids or N-protected (D) α -amino acids of general formula (13), a coupling reagent such as DCC or EDCI and 1-hydroxybenzotriazole hydrate (HOBT) in a solvent such as DMF or methylene chloride to provide amides of general formula (14). Amides of general formula (14) may be deprotected with acid such as 4N HCl in 1,4-dioxane or trifluoroacetic acid in methylene chloride and then treated with 1,1-carbonyldiimidazole (CDI) and a base such as triethylamine or diisopropylamine in a solvent such as acetonitrile to provide 2,5-dioxo-1-imidazolidines of general formula (15). 2,5-Dioxo-1-imidazolidines of general formula (15) may be treated with a base such as sodium hydride and an alkyl halide to provide 2,5-dioxo-1-imidazolidines of general formula (16).

Scheme 5




Alkoxy pyrrolidines of general formula (21), wherein R_2 and R_8 are as defined in formula I, may be prepared as described in Scheme 5. Chlorides of general formula (2), from Scheme 1, may be treated with 3-hydroxypyrrolidine (or (3R)-hydroxypyrrolidine or (3S)-hydroxypyrrolidine), a base such as potassium carbonate, and potassium iodide in a solvent such as 2-butanone with heat to provide hydroxy pyrrolidines of general formula (20). Hydroxy pyrrolidines of general formula (20) may be treated with a base such as sodium hydride and alkyl halides, acid chlorides, carbamyl chlorides, sulfonyl chlorides or chlorophosphates in a solvent such as THF or DMF to provide alkoxy pyrrolidines of general formula (21).

Scheme 6

Alkoxy pyrrolidines of general formula (26), wherein R_2 , R_7 and R_8 are as defined in formula I, may be prepared as described in Scheme 6. Chlorides of general formula (2), from Scheme 1, may be treated with 3-pyrrolidinone, a base such as potassium carbonate, and potassium iodide in a solvent such as 2-butanone with heat to provide ketones of general formula (24). Ketones of general formula (24) may be treated with alkyl halides, magnesium metal and 1,2-dibromoethane (or ketones of general formula (24) may be treated with an alkyl lithium reagent) in a solvent such as THF or diethyl ether to provide tertiary alcohols of general formula (25). Tertiary alcohols of general formula (25) may be treated with a base such as sodium hydride and an alkyl halide in a solvent such as THF or DMF to provide alkoxy pyrrolidines of general formula (26).

Scheme 7



3-Substituted pyrrolidines of general formula (33), wherein R₁, R₇, R₈, and R₉ are as defined in formula I, may be prepared as described in Scheme 7. Phenols of general formula (1), from Scheme 1, may be treated with haloketones of general formula (30), a base such as potassium carbonate, and potassium iodide in a solvent such as 2-butanone with heat to provide ketones of general formula (31). Ketones of general formula (31) may be treated with sodium cyanoborohydride under acidic conditions (or other standard reductive-amination conditions) to provide 3-substituted pyrrolidines of general formula (33).

Schemes 1-7 exemplify preparation of compounds of the present invention wherein Z, as defined in formula I, is CH₂. Compounds of the present invention wherein Z, as defined in formula I, is a covalent bond may be prepared by using the methods described in schemes 1-7 and substituting 1-bromo-2-chloroethane for 1-bromo-3-chloropropane.

The compounds and processes of the present invention will be better understood by reference to the following examples, which are intended as an illustration of and not a limitation upon the scope of the invention. Further, all citations herein are incorporated by reference.

Example 1

N-[(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl]cyclopropanecarboxamide
 1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone (0.05 g, 0.19 mmol) in dichloromethane (2 mL) was treated with cyclopropane carboxylic acid (0.025 g, 0.29 mmol) followed by N-cyclohexylcarbodiimide, N'-methyl polystyrene resin (0.15 g, 0.29 mmol, 2mmol/g loading) and catalytic N,N-dimethylaminopyridine. After shaking at ambient temperature for 14 hours, the mixture was treated with tris-(2-aminoethyl)-amine-polystyrene

(0.063 g, 0.19 mmol, 3 mmol/g loading) resin and the reaction mixture shaken for an additional 2 hours. The mixture was filtered and the filtrate was concentrated. The residue was purified using reverse phase HPLC to afford 0.04 g (63 %) of the title compound.

MS (ESI+) m/z 331 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.2 (d, 1 H), 7.9 (d, 2 H), 5 7.00 (d, 2 H), 4.2 (m, 1 H), 4.15 (t, 2 H), 2.7-2.8 (m, 2 H), 2.5 (m, 2 H), 2.2 (m, 2 H), 1.9 (s, 3 H), 1.8 (m, 2 H), 1.6 (m, 2 H), 0.6 (m, 4 H).

Example 2

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-[4-methyl-2-
10 pyrimidinyl)sulfanyl]acetamide

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and [(4-methyl-2-pyrimidinyl)sulfanyl]acetic acid were processed as described in Example 1 to provide the title compound.

MS (ESI+) m/z 429 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.4 (d, 2 H), 8.2 (d, 1 H), 7.8 15 (d, 2 H), 7.0 (d, 2 H), 6.92 (d, 2 H), 4.15 (m, 3 H), 3.7 (s, 2 H), 2.65-2.7 (m, 2 H), 2.3 (s, 3 H), 2.25 (m, 2 H), 2.0 (m, 2 H), 1.90 (s, 3 H), 1.85 (m, 2 H), 1.6 (m, 2 H).

Example 3

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}nicotinamide

20 1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and nicotinic acid were processed as described in Example 1 to provide the title compound.

MS (ESI+) m/z 368 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 9.0 (s, 1 H), 8.6 (d, 1 H), 8.55 (d, 1 H), 8.1 (d, 1 H), 7.8 (d, 2 H), 7.4 (d, 1 H), 7.0 (d, 2 H), 4.25 (m, 1 H), 4.0 (t, 2 H), 2.8 (t, 2 H), 2.6 (m, 2 H), 2.2 (m, 2 H), 1.95 (m, 2 H), 1.8 (s, 3 H), 1.75 (m, 2 H).

25

Example 4

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-(3-pyridinyl)-1,3-thiazole-4-
carboxamide

30 1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 2-(3-pyridinyl)-1,3-thiazole-4-carboxylic acid were processed as described in Example 1 to provide the title compound.

MS (ESI+) m/z 451 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 9.25 (s, 1 H), 8.7 (d, 1 H), 8.4 (m, 2 H), 7.90 (d, 2 H), 7.6 (m, 2 H), 7.0 (d, 2 H), 4.25 (m, 1 H), 4.2 (t, 2 H), 2.8 (m, 2 H), 2.6 (m, 2 H), 2.2 (m, 2 H), 2.0 (m, 2 H), 1.95 (s, 3 H), 1.8 (m, 2 H).

5

Example 5

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-cyanobenzenesulfonamide
 1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 4-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 6

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-propanesulfonamide
 1-(4-{3-[(3R)-3-aminopyrrolidinyl]propoxy}phenyl)ethanone and 2-propanesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 7

1-(4-{3-[(3R)-3-aminopyrrolidinyl]propoxy}phenyl)-1-propanone
 1-(4-Hydroxyphenyl)-1-propanone was processed as described in Examples 9A-C to provide the title compound.

20 MS (ESI+) m/z 276 (M+H)⁺; ¹H NMR (300 MHz, D₂O) δ 8.0 (d, 2 H), 7.0 (d, 2 H), 4.25 (m, 3 H), 3.8 (br s, 2 H), 3.6 (m, 2 H), 3.0 (q, 2 H), 2.8 (br s, 2 H), 2.2 (m, 2 H), 1.1 (t, 3 H).

Example 8

(5R)-3-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-methyl-2,4-imidazolidinedione
 1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and (2R)-2-[(tert-butoxycarbonyl)amino]propanoic acid were processed as described in Examples 9D-F to provide the title compound.

25 MS (ESI+) m/z 360 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.2 (br s, 1 H), 7.9 (d, 2 H), 7.0 (d, 2 H), 4.4 (m, 1 H), 4.15 (t, 2 H), 4.0 (m, 1 H), 2.85 (m, 2 H), 2.8 (m, 2 H), 2.6 (m, 2 H), 2.0 (m, 2 H), 1.9 (s, 3 H), 1.4 (m, 2 H), 1.2 (d, 3 H).

Example 9

(5S)-3-[(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl]-5-methyl-2,4-imidazolidinedione

Example 9A

5

1-[4-(3-chloropropoxy)phenyl]ethanone

1-(4-hydroxyphenyl)ethanone, 1-bromo-3-chloropropane and potassium carbonate were processed as described in Example 67B to provide the title compound which was used in the next step without further purification.

10

Example 9B

tert-butyl (3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinylcarbamate

1-[4-(3-Chloropropoxy)phenyl]ethanone (1.00 g, 4.7 mmol) in acetone (35 mL), was treated with potassium carbonate (0.97 g, 7.05 mmol), tert-butyl (3R)-pyrrolidinylcarbamate (0.88 g, 4.7 mmol) and potassium iodide (0.78 g, 4.7 mmol). After heating the reaction mixture at reflux for 24 hours, the acetone was evaporated and 50 mL dichloromethane was added. The organic layer was washed with water (3 x 15 mL) and then concentrated under reduced pressure to provide the title compound which was used in the next step without further purification.

20

Example 9C

1-(4-[(3R)-3-aminopyrrolidinyl]propoxy)phenyl)ethanone

tert-Butyl (3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinylcarbamate, from Example 9B, in dichloromethane (25 mL) was treated with trifluoroacetic acid (5 mL). After stirring for 2 hours, the mixture was treated with 2N NaOH. The organic layer was collected, dried over MgSO₄, filtered and concentrated under reduced pressure to provide the title compound as a brown oil (1.2 g).

MS (ESI+) m/z 263 (M+H)⁺; ¹H NMR (300 MHz, D₂O) δ 8.00 (d, 2 H), 7.00 (d, 2 H), 4.2 (t, 2 H), 3.3 (m, 2 H), 3.0 (m, 2 H), 2.5-2.8 (m, 5 H), 2.44 (s, 3 H), 2.1 (m, 2 H); ¹H NMR (300 MHz, DMSO-d₆) δ 8.01 (d, 2H), 7.05 (d, 2H), 4.3 (t, 2H), 3.6 (m, 1H), 2.55 (m, 1H), 2.49 (s, 3H), 2.4-2.3 (m, 4H), 2.2 (m, 1H), 1.8-1.6 (m, 2H), 1.4 (m, 2H).

Example 9Dtert-butyl (1S)-2-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}amino)-1-methyl-2-oxoethylcarbamate

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone (0.1 g, 0.38 mmol) in
5 anhydrous dichloromethane (10 mL), was treated with (2S)-2-[(tert-
butoxycarbonyl)amino]propanoic acid (0.11g, 0.40 mmol), 1-[3-(dimethylamino)propyl]-3-
ethylcarbodiimide hydrochloride (0.077 g, 0.40 mmol) and 1-hydroxybenzotriazole (0.05 g,
0.40 mmol). After stirring at ambient temperature for 12 hours, the mixture was treated with
10 water (20 mL) and the two phases were separated. The aqueous layer was extracted with
dichloromethane (3x15 mL). All the organics were combined, dried over MgSO₄ and
evaporated under reduced pressure to provide the title compound which was used for the
subsequent step without further purification.

Example 9E

15 (2S)-N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-2-aminopropanamide
tert-Butyl (1S)-2-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}amino)-1-
methyl-2-oxoethylcarbamate, from Example 9D, was treated with a 10% solution of TFA in
dichloromethane (30 mL). After stirring for 2 hours, the mixture was treated with water and
the aqueous portion adjusted to pH=8 using 2N NaOH. The organic layer was separated and
20 the aqueous layer extracted with ethyl acetate (3x20 mL). All the organic layers were
combined, dried over MgSO₄, filtered and evaporated under reduced pressure to provide the
title compound.

Example 9F

25 (5S)-3-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-methyl-2,4-imidazolidinedione
(2S)-N-((3R)-1-[3-(4-Acetylphenoxy)propyl]pyrrolidinyl)-2-aminopropanamide in
acetonitrile (25 mL) was treated with 1,1'-carbonyldiimidazole (0.05 g, 0.34 mmol) and
triethylamine (0.07 g, 0.68 mmol). After heating at 40 °C for 16 hours, the mixture was
allowed to cool to ambient temperature and the solvent was evaporated under reduced
30 pressure. The residue was purified via reverse-phase silica gel chromatography using a C18
column, to provide 0.07 g (49 %) of the title compound.

MS (ESI+) m/z 360 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.2 (br s, 1 H), 7.9 (d, 2 H), 7.0 (d, 2 H), 4.4 (m, 1 H), 4.15 (t, 2 H), 3.95 (m, 1H), 2.8 (m, 3 H), 2.51 (m, 4 H), 2.1 (m, 2 H), 1.8 (m, 2 H), 1.4 (m, 2 H), 1.1 (d, 3 H).

5

Example 10(5R)-3-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-isopropyl-2,4-imidazolidinedione

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and (2S)-2-[(tert-butoxycarbonyl)amino]-3-methylbutanoic acid were processed as described in Examples 9D-F to provide the title compound.

10

Example 114-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

15

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 12N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-propanesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-propanesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 13N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-(3-pyridinyl)-1,3-thiazole-4-carboxylic acid were processed as described in Example 1 to provide the title compound.

30

Example 14N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-4-cyanobenzamide

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 4-cyanobenzoic acid were processed as described in Example 1 to provide the title compound.

5 MS (ESI+) m/z 392 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.00 (d, 2 H), 7.8-7.9 (m, 4 H), 7.00 (d, 2 H), 4.4 (m, 1 H), 4.15 (m, 2 H), 2.8 (m, 2 H), 2.5-2.7 (m, 4 H), 2.2 (m, 2 H), 1.95 (s, 3 H), 1.8 (m, 2 H).

Example 15N-((3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl)pyrrolidinyl)benzenesulfonamide

10 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 164-bromo-N-((3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl)pyrrolidinyl)benzenesulfonamide

20 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-bromobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 17N-((3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl)pyrrolidinyl)-4-fluorobenzenesulfonamide

25 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 184-chloro-N-((3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl)pyrrolidinyl)benzenesulfonamide

30

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 19

N-(4-{{[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl]amino}sulfonyl}phenyl)acetamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(acetylamino)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 20

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-methoxybenzenesulfonamide

15 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (0.05 g, 0.15 mmol) in dichloromethane (2 mL) was treated with polymer supported N,N-diisopropylethylamine (0.10 g, 0.31 mmol, 3mmol/g loading), catalytic N,N-dimethylaminopyridine, and 4-methoxybenzenesulfonyl chloride (0.034 g, 0.17 mmol). After shaking at ambient temperature for 14 hours, the mixture was treated with tris-(2-aminoethyl)-amine-polystyrene (0.05 g, 0.15 mmol, 3 mmol/g loading) resin and the reaction mixture shaken for an additional 2 hours. The reaction mixture was filtered and the filtrate was concentrated. The residue was purified using reverse phase HPLC to afford 0.053 g (79%) of the title compound.

20 25 MS (ESI+) m/z 492 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 7.85 (m, 4 H), 7.7 (m, 4 H), 7.0 (m, 4 H), 4.0 (t, 2 H), 3.8 (s, 3 H), 3.6 (m, 1 H), 2.4-2.6 (m, 4 H), 2.2 (m, 1 H), 1.8 (m, 4 H), 1.5 (m, 1 H).

30

Example 21

4-tert-butyl-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-tert-butylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 22

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 23

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(trifluoromethyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 24

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2,5-dimethoxybenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2,5-dimethoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 25

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-methylbenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

MS (ESI+) m/z 476 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 7.8 (m, 4 H), 7.7 (d, 2 H), 7.4 (m, 4 H), 7.0 (d, 2 H), 4.4 (s, 3 H), 4.0 (t, 2 H), 3.7 (m, 1 H), 2.8 (m, 1 H), 2.5 (m, 4 H), 2.2 (m, 1 H), 2.0 (m, 1 H), 1.9 (m, 2 H), 1.6 (m, 1 H).

5

Example 26

3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-chloro-4-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 27

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-ethylbenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-ethylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 28

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(2-propyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 29

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 30

2-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

5 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 31

10 3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 32

3,5-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

20 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3,5-dichlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 33

4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

25 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 34

3-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

MS (ESI+) m/z 487 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.2 (s, 1 H), 8.1 (d, 2 H), 7.8 (m, 5 H), 7.6 (d, 2 H), 7.0 (d, 2 H), 4.0 (t, 2 H), 3.7 (m, 1 H), 2.8 (m, 1 H), 2.4-2.6 (m, 5 H), 2.2 (m, 1 H), 1.8-2.0 (m, 2 H), 1.5 (m, 1 H).

10

Example 35

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 1-methyl-1H-imidazole-4-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 36

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-thiophenesulfonamide

20

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-thiophenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 37

5-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-methyl-1-benzothiophene-2-sulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 5-chloro-3-methyl-1-benzothiophene-2-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 38

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3,5-bis(trifluoromethyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 39

N-(5-{|((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)amino]sulfonyl}-4-methyl-1,3-thiazol-2-yl)acetamide

10 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-(acetylamino)-4-methyl-1,3-thiazole-5-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 40

15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-8-quinolinesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 8-quinolinesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 41

4-butoxy-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

25 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-butoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 42

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide

30

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3,4-dimethoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 43

3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-chloro-4-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 44

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-phenylethanesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-phenylethanesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 45

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-(trifluoromethoxy)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 46

2-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

30

Example 47N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-3-methylbenzenesulfonamide

5 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 48N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide

10 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(trifluoromethyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 49N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-3-fluorobenzenesulfonamide

20 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 50N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide

25 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(trifluoromethoxy)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 51

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2,4-difluorobenzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 2,4-difluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 52

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-7-isoquinolinesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 7-isoquinolinesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 53

N-(2-chloro-4-{[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl]amino}sulfonyl)phenyl)acetamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 4-(acetylamino)-3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 54

3,4-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and 3,4-dichlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 55

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 564-bromo-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-bromobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 57N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide

15

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 584-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 59N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methoxybenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-methoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 60

5 4-tert-butyl-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide
(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-tert-butylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 61

10 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide
(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 62

20 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-(trifluoromethyl)benzenesulfonamide
(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-trifluoromethylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 63

25 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2,5-dimethoxybenzenesulfonamide
(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2,5-dimethoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 64 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-methylbenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 653-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-chloro-4-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 66N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-ethylbenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-ethylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 67

20 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide

Example 67Acyclopropyl(4-hydroxyphenyl)methanone

Sodium hydroxide in water (50% (w/w), 40.4 mL) was treated over a period of 15 minutes with para-hydroxy-4-chlorobutyrophenone (25.0 g, 137 mmol), followed by additional aqueous sodium hydroxide (25% (w/w), 177 mL). Additional para-hydroxy-4-chlorobutyrophenone (25.0 g, 137 mmol) was added portionwise to the reaction mixture, followed by solid sodium hydroxide (40.4 g) resulting in the formation of a yellow precipitate. After refluxing for 60 minutes, water (50 mL) was added and the resulting mixture was refluxed for another 60 minutes, allowed to cool to ambient temperature, diluted with water (100 mL) and neutralized with acetic acid. The precipitate was collected by

filtration, washed with water, air-dried, and triturated at 40 °C with chloroform (1.5 L). The chloroform solution was dried (MgSO_4), filtered, and concentrated. The residue was crystallized from chloroform/hexanes to provide 26.65 g (95%) of the title compound.

MS (APCI(-)) m/z 161 (M-H)⁺; ¹H NMR (300 MHz, CDCl_3) δ 7.9 (d, 2H), 6.9 (d, 2H), 2.65 (m, 1H), 1.23 (m, 2H), 1.03 (m, 2H).

Example 67B

(4-(3-chloropropoxy)phenyl)(cyclopropyl)methanone

Cyclopropyl(4-hydroxyphenyl)methanone (10 g, 61.7 mmol), K_2CO_3 (12.7 g, 91.9 mmol), and 1-bromo-3-chloropropane (10.74 g, 68.2 mmol) in 2-butanone (100 mL) were refluxed for 24 hours, allowed to cool to ambient temperature, filtered, and concentrated. The residue was heated at 40 °C under reduced pressure for three hours to provide the title compound of sufficient purity for subsequent use without further purification (13.25 g, 90%).

Example 67C

15 (4-{3-[{(3R)-3-aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone

(4-(3-Chloropropoxy)phenyl)(cyclopropyl)methanone was processed as described in Examples 9B and 9C to provide the title compound.

Example 67D

20 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide

(4-{3-[{(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-(2-propyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25 MS (ESI+) m/z 472 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d_6) δ 8.0 (d, 2 H), 7.7 (d, 2 H), 7.4 (d, 2 H), 7.0 (d, 2 H), 4.1 (t, 2 H), 3.8 (br s, 1 H), 3.0 (m, 1 H), 2.85 (m, 1 H), 2.48-2.6 (m, 4 H), 2.2 (m, 1 H), 1.8-1.95 (m, 2 H), 1.5 (m, 1 H), 1.1 (d, 6 H), 1.0 (d, 4 H).

Example 68

30 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 692-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 703-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

20

Example 713-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

30

Example 72N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-3-fluorobenzenesulfonamide

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 3-

fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 73N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-(trifluoromethoxy)benzenesulfonamide

1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 4-(trifluoromethoxy)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

MS (ESI+) m/z 486 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d6) δ 8.0 (d, 2 H), 7.7 (d, 2 H), 7.6 (d, 2 H), 7.0 (d, 2 H), 4.1 (t, 2 H), 3.6 (m, 1 H), 2.8 (m, 2 H), 2.4-2.5 (m, 2 H), 2.2 (m, 1 H), 1.8-1.9 (m, 4 H), 1.45 (m, 1 H), 1.25 (s, 9 H), 1.00 (d, 4 H).

10

Example 74N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2,4-difluorobenzenesulfonamide
1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 2,4-difluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 75N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-5-isoquinolinesulfonamide
1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 5-isoquinolinesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 76N-{4-[(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl]amino}sulfonyl]-2-chlorophenyl}acetamide
1-(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 4-(acetylamino)-3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 77N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-3,4-dichlorobenzenesulfonamide

1-(4-{3-[3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 3,4-dichlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 78

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1H-pyrrole-2-sulfonamide

1-(4-{3-[3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1H-pyrrole-2-sulfonyl chloride were processed as described in 10 Example 20 to provide the title compound.

Example 79

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]oxy}benzenesulfonamide

15 1-(4-{3-[3R)-3-Aminopyrrolidinyl]propoxy}phenyl)ethanone and 4-{[3-chloro-5-(trifluoromethyl)-2-pyridinyl]oxy}benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 80

20 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide

(4-{3-[3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 1-methyl-1H-imidazole-4-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

Example 81

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-thiophenesulfonamide

30 (4-{3-[3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-thiophenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 825-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-methyl-1-benzothiophene-2-sulfonamide

5 (4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 5-chloro-3-methyl-1-benzothiophene-2-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 83

10 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3,5-bis(trifluoromethyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 84

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-quinolinesulfonamide

20 (4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 8-quinolinesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 85

25 4-butoxy-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-butoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30 MS (ESI+) m/z 501 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.0 (d, 2 H), 7.65 (d, 2 H), 7.6 (d, 1 H), 7.15 (d, 2 H), 7.0 (d, 2 H), 4.1 (m, 4 H), 3.6 (m, 1 H), 2.8 (m, 1 H), 2.4-2.7 (m, 4 H), 2.2 (m, 1 H), 1.65-1.8 (m, 6 H), 1.4 (m, 3 H), 1.00 (d, 4 H), 0.95 (t, 3 H).

Example 86N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide

5 ((4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3,4-dimethoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 87

10 3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-chloro-4-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

Example 88N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-phenylethenesulfonamide

20 ((4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-phenylethenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 89N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-(trifluoromethoxy)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

30

Example 90

2-cyano-N-((3R)-1-{3-[4-

(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 2-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the

5 title compound.

Example 91

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-
methylbenzenesulfonamide

10 (4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-methylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 92

15 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-
(trifluoromethyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-trifluoromethylbenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 93

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-
fluorobenzenesulfonamide

25 (4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3-fluorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 94

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-
(trifluoromethoxy)benzenesulfonamide

30

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-(trifluoromethoxy)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 95

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-isoquinolinesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 8-isoquinolinesulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 96

N-(2-chloro-4-{[(3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl]amino}sulfonyl)phenyl)acetamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 4-(acetylamino)-3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

15

20

Example 97

3,4-dichloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}phenyl)(cyclopropyl)methanone and 3,4-dichlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

25

30

Example 98

tert-butyl 1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}-3-pyrrolidinylcarbamate

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile and tert-butyl pyrrolidinylcarbamate were processed as described in Examples 9A-9C to provide the title compound.

Example 994'-{3-[(3R)-3-aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile

4'-Hydroxy[1,1'-biphenyl]-4-carbonitrile was processed as described in Examples 9A-C to provide the title compound.

5 MS (APCI+) m/z 322 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.25 (m, 1H), 1.80 (m, 1H), 1.99-2.20 (m, 2H), 2.74-3.09 (m, 4H), 3.20 (m, 1H), 3.78 (m, 1H), 4.08 (t, 2H, J=6.0 Hz), 4.39-5.03 (m, 2H), 6.97 (d, 2H, J=8.8 Hz), 7.52 (d, 2H, J=8.5 Hz), 7.62 (d, 2H, J=8.5 Hz), 7.69 (d, 2H, J=8.5 Hz).

10

Example 100tert-butyl (3S)-1-{2-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]ethyl}pyrrolidinylcarbamate

4'-Hydroxy[1,1'-biphenyl]-4-carbonitrile, 1-bromo-2-chloroethane and tert-butyl (3S)-pyrrolidinylcarbamate were processed as described in Examples 9A and 9B to provide the title compound.

15

Example 1014-methoxy-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 4-methoxybenzenesulfonyl chloride were processed as described in Example 20 to provide the 20 title compound.

Example 1024-isopropyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

25

Example 102A(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine

4-(2-Pyridinyl)phenol was processed as described in Examples 9A-9C to provide the title compound.

30 MS (ESI+) m/z 298 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.45 (m, 1H), 7.75 (d, 2H), 7.6 (m, 1H), 7.55 (m, 1H), 7.1 (m, 1H), 6.9 (d, 2H), 4.25 (t, 2H), 3.7 (m, 1H), 2.5-2.3 (m, 5H), 2.10 (m, 1H), 1.8-1.6 (m, 2H), 1.4 (m, 2H).

Example 102B4-isopropyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 4-(2-propyl)benzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

MS (ESI+) m/z 480 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.66 (d, 1H), 7.95 (m, 2H), 7.85 (d, 1H), 7.70 (m, 3H), 7.37 (d, 1H), 7.20 (m, 1H), 6.94 (m, 3H), 4.1 (t, 2H), 3.8 (br s, 1H), 3.0 (m, 1H), 2.85 (m, 1H), 2.48-2.6 (m, 4H), 2.2 (m, 1H), 1.8-1.95 (m, 2H), 1.5 (m, 1H), 1.1 (d, 6H).

Example 1034-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 4-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

Example 1043-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 3-cyanobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

MS (ESI+) m/z 463 (M+H)⁺; ¹H NMR (300 MHz, DMSO-d₆) δ 8.56 (d, 1 H), 8.21 (d, 1 H), 8.18 (s, 1 H), 7.88 (d, 2 H), 7.72 (m, 1 H), 7.5 (m, 2 H), 7.45 (m, 1 H), 7.0 (m, 1 H), 6.85 (d, 2 H), 4.1 (t, 2 H), 3.6 (m, 1 H), 2.8 (m, 2 H), 2.4-2.5 (m, 2 H), 2.2 (m, 1 H), 1.8-1.9 (m, 4 H), 1.45 (m, 1 H).

Example 1051-methyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)-1H-imidazole-4-

30 sulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 1-methyl-1H-imidazole-4-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

5

Example 1063,4-dimethoxy-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 1-methyl-1H-imidazole-4-sulfonyl chloride were processed as described in Example 20 to provide the title compound.

10

Example 107N-(2-chloro-4-{[(3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)amino]sulfonyl}phenyl)acetamide

15

(3R)-1-{3-[4-(2-Pyridinyl)phenoxy]propyl}pyrrolidinylamine and 4-(acetylamino)-3-chlorobenzenesulfonyl chloride were processed as described in Example 20 to provide the title compound.

20

Example 1084'-{3-[(3R)-3-(dimethylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (200 mg, 0.74 mmol), N,N-dimethyl-N-[(3R)-pyrrolidinyl]amine (85 mg, 0.74 mmol), 250 mg of potassium carbonate and 300 mg of potassium iodide in 20 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound.

25

MS (ESI+) m/z 350 (M+H)⁺; ¹³C NMR(500 MHz, CD₃OD) 29.3, 29.6, 43.9, 54.2, 54.3, 59.6, 66.5, 67.3, 111.0, 116.2, 119.9, 128.2, 129.4, 132.7, 133.7, 146.7, 161.2; ¹H NMR (500 MHz, CD₃OD) 1.74 (m, 1H), 2.0 (m, 2H), 2.02 (m, 1H) 2.23 (s, 6H), 2.32 (m, 1H), 2.51 (m, 1H), 2.62 (m, 1H), 2.71 (m, 1H), 2.84 (m, 2H), 2.97 (m, 1H), 4.08 (t, J=7 Hz, 2H), 7.02 (d, J=11 Hz, 2H), 7.61 (d, J=11 Hz, 2H), 7.74 (s, 4H).

30

Example 109tert-butyl (3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl(methyl)carbamate

5 4'-{3-[{3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (2.31 g, 8.54 mmol), tert-butyl methyl[(3R)-pyrrolidinyl]carbamate (1.71 g, 8.54 mmol), potassium carbonate (2.95 g, 21.34 mmol) and potassium iodide (3.54 g, 21.34 mmol) in 75 mL of 2-butanone were heated at 115 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (89% yield).

10 MS (APCI+) m/z 436 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.45 (s, 9H), 1.8 (m, 2H), 2.1 (m, 2H), 2.4 (m, 2H), 2.7 (m, 2H), 2.85 (s, 3H), 4.1 (t, J=7Hz, 2H), 4.25 (m, 1H), 4.75 (m, 1H), 6.98 (d, J=9Hz, 2H), 7.5 (d, J=9Hz, 2H), 7.65 (m, 4H).

15

Example 1104'-{3-[{3R)-3-(methylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile

The product from Example 109 (3.3 g, 7.58 mmol) in dichloromethane (100 mL) was treated with trifluoroacetic acid (6.0 ml) at 0 °C. After stirring at ambient temperature for 18 hours, the mixture was evaporated under reduced pressure. The residue was partitioned between saturated aqueous sodium bicarbonate (2 x 30 mL) and dichloromethane (150 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate and filtered. The filtrate was evaporated under reduced pressure and the residue purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (95% yield).
MS (APCI+) m/z 336 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.62 (m, 1H), 2.00 (m, 2H), 2.13 (m, 1H), 2.26 (m, 1H), 2.41 (s, 3H), 2.43-2.83 (m, 5H), 3.25 (m, 1H), 4.08 (t, 2H, J=6.3 Hz), 6.99 (d, 2H, J=8.9 Hz), 7.52 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=16.8, 8.6 Hz).

20

25

Example 111N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-N-methylacetamide

30 The product from Example 110 (50 mg, 0.15 mmol), acetic anhydride (0.021 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane

were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (60% yield).

MS (APCI+) m/z 378 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.12 (m, 5H), 2.32 (m, 7H), 2.65 (s, 3H), 2.83-3.18 (m, 7H), 3.23-3.79 (m, 9H), 3.87-4.20 (m, 7H), 5.38 (m, 1H), 5.64-5.94 (m, 1H), 6.97 (m, 2H), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.2, 8.6 Hz).

Example 112

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,3,3-trimethylbutanamide

The product from Example 110 (50 mg, 0.15 mmol), tert-butylacetyl chloride (0.031 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 434 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.05 (s, 9H), 2.11-2.41 (m, 5H), 2.90 (m, 1H), 3.08 (m, 3H), 3.24-3.47 (m, 3H), 3.57-3.80 (m, 2H), 3.94 (m, 1H), 4.12 (m, 3H), 6.97 (m, 2H), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.3, 8.8 Hz).

Example 113

methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl(methyl)carbamate

The product from Example 110 (50 mg, 0.15 mmol), dimethyl dicarbonate (0.024 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.3 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃/MeOH/NH₄OH, 9:1:0.1) to provide the title compound (60% yield).

MS (APCI+) m/z 394 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 0.89 (m, 2H), 1.30-1.52 (m, 2H), 1.44 (s, 3H), 1.77 (m, 1H), 2.15-2.38 (m, 3H), 2.65 (s, 3H), 2.91 (m, 3H), 3.18-3.52 (m, 3H), 3.73 (m, 1H), 3.84-4.04 (m, 1H), 4.05-4.26 (m, 2H), 6.97 (d, 2H, J=8.5 Hz), 7.53 (d, 2H, J=8.5 Hz), 7.66 (dd, 4H, J=20.4, 8.1 Hz).

Example 114tert-pentyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl(methyl)carbamate

5 The product from Example 110 (50 mg, 0.15 mmol), di(tert-pentyl) dicarbonate (0.055 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.3 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (60% yield).

10 MS (APCI+) m/z 450 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 0.90 (s, 2H), 1.45 (m, 4H), 1.80 (m, 2H), 2.3 (m, 2H), 2.5 (m, 1H), 2.7 (s, 2H), 2.85 (s, 2H), 2.95 (s, 3H), 3.4 (m, 4H), 3.75 (m, 1H), 3.95 (m, 1H), 4.15 (t, J=7Hz, 2H), 5.05 (m, 1H), 6.98 (d, J=9Hz, 2H), 7.55 (d, J=9Hz, 2H), 7.65 (m, 4H).

15

Example 115N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N',N'-trimethylurea

The product from Example 110 (50 mg, 0.15 mmol), dimethylcarbamyl chloride (0.021 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

15 ¹H NMR (300 MHz, CDCl₃) δ 2.14-2.40 (m, 3H), 2.64 (s, 3H), 2.84 (s, 3H), 2.85 (s, 3H), 3.25-3.46 (m, 3H), 3.52 (m, 1H), 3.61-3.81 (m, 1H), 3.98 (m, 1H), 4.13 (m, 2H), 5.73-6.09 (m, 1H), 6.98 (m, 2H), 6.98 (m, 2H), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.1, 8.6 Hz).

25 MS (APCI+) m/z 407 (M+H)⁺.

Example 116N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-1-pyrrolidinecarboxamide

30 The product from Example 110 (50 mg, 0.15 mmol), 1-pyrrolidinecarbonyl chloride (0.030 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of

dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 433 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.85 (m, 3H), 2.17-2.37 (m, 3H), 2.64 (m, 1H), 2.89 (m, 3H), 3.23-3.46 (m, 6H), 3.62-3.83 (m, 1H), 3.95 (m, 1H), 4.12 (m, 2H), 6.98 (dd, 2H, J=11.4, 8.6 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 1H, J=20.2, 8.6 Hz).

Example 117

N'-tert-butyl-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylurea

The product from Example 110 (50 mg, 0.15 mmol), tert-butylcarbamyl chloride (0.026 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 435 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.35 (s, 9H), 2.15-2.44 (m, 4H), 2.85 (s, 3H), 3.10-3.43 (m, 3H), 3.57-3.78 (m, 1H), 3.81-4.01 (m, 1H), 4.13 (m, 2H), 4.23-4.50 (m, 1H), 5.08-5.37 (m, 0H), 6.98 (m, 2H), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.3, 8.5 Hz).

20

Example 118

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-morpholinecarboxamide

The product from Example 110 (50 mg, 0.15 mmol), 4-morpholinecarbonyl chloride (0.026 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 449 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.14-2.39 (m, 3H), 2.89 (d, 3H, J=16.9 Hz), 3.12-3.56 (m, 7H), 3.61-3.82 (m, 5H), 3.87-4.22 (m, 3H), 4.42-4.65 (m, 1H), 4.70-5.24 (m, 2H), 6.97 (m, 2H), 7.53 (d, 2H, J=8.5 Hz), 7.66 (dd, 4H, J=20.3, 8.4 Hz).

5

Example 119

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-methylbenzamide

The product from Example 110 (50 mg, 0.15 mmol), 4-fluorobenzoyl chloride (0.026 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

10 MS (APCI+) m/z 458 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.22-2.58 (m, 3H), 2.64 (s, 3H), 2.93-3.19 (m, 3H), 3.27-3.94 (m, 5H), 3.96-4.22 (m, 3H), 4.23-4.58 (m, 1H), 6.98 (d, 2H, J=8.4 Hz), 7.13 (t, 2H, J=8.5 Hz), 7.44 (dd, 2H, J=8.8, 5.5 Hz), 7.53 (d, 2H, J=8.9 Hz), 7.66 (dd, 4H, J=20.7, 8.8 Hz).

Example 120

4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylbenzamide

20 The product from Example 110 (50 mg, 0.15 mmol), 4-cyanobenzoyl chloride (37 mg, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

25 MS (APCI+) m/z 465 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.18-2.56 (m, 4H), 2.63 (s, 3H), 2.89-3.17 (m, 4H), 3.29-3.60 (m, 4H), 3.72-3.96 (m, 1H), 4.03-4.24 (m, 3H), 6.97 (d, 2H, J=8.8 Hz), 7.53 (d, 4H, J=7.1 Hz), 7.66 (dd, 4H, J=20.9, 8.2 Hz), 7.75 (d, 2H, J=8.2 Hz).

30

Example 121

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylnicotinamide

The product from Example 110 (50 mg, 0.15 mmol), nicotinoyl chloride (40 mg, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 441 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.20-2.61 (m, 3H), 2.66 (s, 2H), 2.91-3.24 (m, 3H), 3.24-3.76 (m, 4H), 3.77-3.97 (m, 1H), 3.99-4.27 (m, 3H), 6.98 (d, 2H, J=8.5 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.1, 8.2 Hz), 7.86-8.07 (m, 1H), 8.08-8.50 (m, 1H), 8.78 (d, 2H, J=14.6 Hz).

10

Example 122

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-furamide

The product from Example 110 (50 mg, 0.15 mmol), 2-furoyl chloride (0.022 mL, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 430 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.19-2.60 (m, 4H), 2.64 (s, 3H), 3.19-3.61 (m, 6H), 3.62-3.88 (m, 2H), 3.90-4.21 (m, 4H), 6.53 (dd, 1H, J=3.4, 1.7 Hz), 6.98 (d, 2H, J=8.1 Hz), 7.06 (m, 1H), 7.53 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=20.0, 8.8 Hz).

Example 123

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide

The product from Example 110 (50 mg, 0.15 mmol), 2-(3-pyridinyl)-1,3-thiazole-4-carbonyl chloride (46 mg, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.3 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 524 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.20 (m, 2H), 2.35 (m, 4H), 2.65 (s, 3H), 3.10 (m, 2H), 3.40 (m, 4H), 4.15 (m, 2H), 6.98 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.65 (m, 4H), 8.05 (s, 1H), 8.40 (m, 1H), 8.70 (m, 1H), 9.20 (m, 1H).

5

Example 124

N-[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl]-N,N',N'-trimethylsulfamide

The product from Example 110 (50 mg, 0.15 mmol), dimethylsulfamoyl chloride (0.024 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

10 MS (APCI+) m/z 443 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.15-2.56 (m, 4H), 2.65 (s, 3H), 2.71-2.92 (m, 2H), 2.83 (s, 6H), 2.97 (m, 1H), 3.18 (m, 1H), 3.36 (m, 2H), 3.85 (m, 1H), 4.12 (m, 2H), 6.97 (d, 2H, J=8.9 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=20.3, 8.4 Hz).

Example 125

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-fluoro-N-methylbenzenesulfonamide

20 The product from Example 110 (50 mg, 0.15 mmol), 3-fluorobenzenesulfonyl chloride (0.039 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).

25 MS (APCI+) m/z 494 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.96-2.43 (m, 3H), 2.64 (s, 2H), 2.73-2.94 (m, 3H), 3.07 (m, 1H), 3.21-3.43 (m, 2H), 4.10 (m, 2H), 6.95 (d, 2H, J=8.8 Hz), 7.34 (m, 1H), 7.53 (d, 2H, J=8.8 Hz), 7.58 (m, 1H), 7.66 (dd, 4H, J=21.2, 8.3 Hz).

Example 126

30 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylbenzenesulfonamide

The product from Example 110 (50 mg, 0.15 mmol), 4-cyanobenzenesulfonyl chloride (45 mg, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 5 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 501 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.30 (m, 2H), 2.65 (s, 3H), 2.77-3.01 (m, 2H), 3.09 (m, 0H), 3.23-3.43 (m, 2H), 4.11 (m, 1H), 6.95 (d, 2H, J=8.8 Hz), 7.54 (d, 2H, J=8.9 Hz), 7.67 (dd, 4H, J=21.2, 8.3 Hz), 7.89 (dd, 4H, J=23.9, 8.3 Hz).

10

Example 127

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-isopropyl-N-methylbenzenesulfonamide

The product from Example 110 (50 mg, 0.15 mmol), 4-(2-propyl)benzenesulfonyl chloride (0.044 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 15 9:1:0.1) to provide the title compound (62% yield).

MS (APCI+) m/z 518 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.27 (d, 6H, J=6.7 Hz), 2.11-2.40 (m, 3H), 2.64 (s, 3H), 2.79 (m, 3H), 2.84-3.14 (m, 2H), 3.20-3.43 (m, 2H), 4.10 (m, 2H), 6.95 (d, 2H, J=8.8 Hz), 7.38 (d, 2H, J=8.2 Hz), 7.52 (d, 2H, J=8.5 Hz), 7.66 (dd, 4H, J=21.4, 20 8.5 Hz), 7.70 (d, 2H, J=8.2 Hz).

Example 128

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-(methylsulfonyl)benzenesulfonamide

The product from Example 110 (50 mg, 0.15 mmol), 4-(methylsulfonyl)benzenesulfonyl chloride (57 mg, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.3 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 30 9:1:0.1) to provide the title compound (68% yield).

MS (APCI+) m/z 554 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.30 (m, 4H), 2.60 (s, 3H), 2.90 (s, 2H), 3.10 (s, 3H), 3.30 (m, 2H), 4.10 (t, J=7Hz, 2H), 4.95 (m, 1H), 6.95 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.65 (m, 4H), 8.00 (d, J=9Hz, 2H), 8.15 (d, J=9Hz, 2H).

5

Example 129N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-(4-fluorobenzoyl)benzamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-fluorobenzoyl chloride (0.028 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (20% yield).

MS (APCI+) m/z 436 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.24-2.56 (m, 4H), 2.56-2.84 (m, 1H), 3.11-3.45 (m, 1H), 3.45-3.78 (m, 4H), 3.93 (m, 1H), 4.12 (m, 2H), 4.27 (m, 1H), 5.35-5.62 (m, 1H), 6.82-7.04 (m, 6H), 7.44 (m, 4H), 7.52 (d, 2H, J=8.8 Hz), 7.65 (dd, 4H, J=20.0, 8.2 Hz).

Example 130N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)acetamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), acetyl chloride (22.0 μL, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (70% yield).

MS (APCI+) m/z 364 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.0 (m, 3H), 2.2 - 2.6 (m, 4H), 2.9 – 3.2 (m, 2H), 3.35 (m, 2H), 3.75 (m, 1H), 4.0 (m, 1H), 4.15 (m, 2H), 4.9 (m, 1H), 6.95 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.65 (m, 4H), 8.3 (m, 2H).

Example 131

30 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,3-dimethylbutanamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), tert-butylacetyl chloride (0.033 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (52% yield).

MS (APCI+) m/z 420 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.02 (s, 9H), 2.10 (s, 2H), 2.21 (m, 1H), 2.35 (m, 2H), 2.42-2.81 (m, 6H), 2.83-3.17 (m, 2H), 3.35 (m, 2H), 3.72 (m, 1H), 3.97 (m, 1H), 4.13 (m, 2H), 4.92 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.54 (d, 2H, J=8.5 Hz), 7.67 (dd, 4H, J=21.8, 8.3 Hz).

10

Example 132

allyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), allyl chloroformate (0.039 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (51% yield).

MS (APCI+) m/z 406 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.17-2.41 (m, 4H), 2.41-2.64 (m, 6H), 2.91 (m, 1H), 3.10 (m, 1H), 3.23-3.46 (m, 2H), 3.81 (m, 2H), 4.00 (m, 1H), 4.14 (m, 2H), 4.56 (m, 2H), 4.68 (m, 1H), 5.20 (dd, 1H, J=10.3, 0.8 Hz), 5.30 (dd, 1H, J=17.3, 1.4 Hz), 5.90 (ddd, 1H, J=22.8, 10.7, 5.7 Hz), 6.80 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=8.4 Hz), 7.67 (dd, 4H, J=21.6, 8.7 Hz).

Example 133

methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), methyl chloroformate (0.025 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (51% yield).

30

MS (APCI+) m/z 380 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.05 (m, 6H), 2.29 (m, 3H), 2.53 (m, 1H), 3.08 (m, 1H), 3.33 (m, 2H), 3.67 (s, 3H), 3.77 (m, 1H), 3.98 (m, 1H), 4.14 (t, 2H, J=5.6 Hz), 4.67 (m, 1H), 6.96 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=9.2 Hz), 7.67 (dd, 4H, J=21.6, 8.6 Hz).

5

Example 134

tert-pentyl (3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinylcarbamate
 4'-{3-[{(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 1,1-dimethylpropyl chloroformate (57.1 μL, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (72% yield).

MS (APCI+) m/z 436 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 0.90 (t, J=7Hz, 3H), 1.4 (s, 6H), 1.75 (m, 2H), 2.2 (m, 1H), 2.35 (m, 2H), 2.5 (m, 1H), 2.9 (m, 4H), 3.35 (m, 2H), 3.7 (m, 1H), 4.0 (m, 1H), 4.15 (m, 2H), 6.95 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.65 (m, 4H).

Example 135

N'-(3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-N,N-dimethylurea
 4'-{3-[{(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), dimethylcarbamyl chloride (0.021 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (65% yield).

MS (APCI+) m/z 393 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.20-2.42 (m, 4H), 2.48-2.77 (m, 6H), 2.90 (m, 1H), 2.93 (s, 6H), 3.07 (m, 1H), 3.34 (m, 2H), 3.76 (m, 1H), 3.96 (m, 1H), 4.14 (t, 2H, J=5.4 Hz), 4.86 (m, 1H), 6.71 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=21.8, 8.7 Hz).

N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-1-pyrrolidinecarboxamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 1-pyrrolidinecarbonyl chloride (0.031 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (68% yield).

MS (APCI+) m/z 419 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.90 (m, 4H), 2.21-2.43 (m, 3H), 2.59 (m, 1H), 2.83-3.50 (m, 12H), 3.77 (m, 1H), 3.97 (m, 1H), 4.15 (t, 2H, J=4.8 Hz), 4.89 (m, 1H), 6.59 (m, 1H), 6.97 (d, 2H, J=8.4 Hz), 7.53 (d, 2H, J=8.5 Hz), 7.67 (dd, 4H, J=21.7, 8.5 Hz).

Example 137

15 N-(tert-butyl)-N'-(3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)urea
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), tert-butylcarbamyl chloride (0.027 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (65% yield).
 20 MS (APCI+) m/z 421 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.31 (s, 9H), 2.16 (m, 1H), 2.35 (m, 2H), 2.50 (m, 1H), 2.61-3.08 (m, 4H), 3.33 (m, 2H), 3.68 (m, 1H), 3.94 (m, 1H), 4.15 (t, 2H, J=5.6 Hz), 4.74 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.54 (d, 2H, J=9.2 Hz), 7.67 (dd, 4H, J=21.7, 8.8 Hz).

25

Example 138

N-((3R)-1-{3-[{4'-cyano[1,1'-biphenyl]-4-yl}oxy]propyl}pyrrolidinyl)-4-morpholinecarboxamide
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-morpholinecarbonyl chloride (0.027 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C

for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (52% yield).

MS (APCI+) m/z 435 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.24 (m, 1H), 2.36 (m, 2H), 2.60 (m, 1H), 2.70-3.01 (m, 4H), 3.10 (m, 1H), 3.33 (m, 2H), 3.43 (t, 4H, J=4.6 Hz), 3.67 (t, 4H, J=4.4 Hz), 3.77 (m, 1H), 3.97 (m, 1H), 4.15 (t, 2H, J=5.1 Hz), 4.88 (m, 1H), 6.97 (d, 2H, J=8.9 Hz), 7.54 (d, 2H, J=8.5 Hz), 7.67 (dd, 4H, J=22.1, 8.5 Hz).

Example 139

10 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluorobenzamide
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-fluorobenzoyl chloride (0.027 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (58% yield).
 15 MS (APCI+) m/z 444 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.36 (m, 3H), 2.58-2.88 (m, 3H), 2.99 (m, 1H), 3.18 (m, 1H), 3.29-3.50 (m, 2H), 3.90 (m, 1H), 4.03 (m, 1H), 4.16 (t, 2H, J=4.8 Hz), 5.16 (m, 1H), 6.96 (d, 2H, J=8.8 Hz), 7.11 (t, 2H, J=8.7 Hz), 7.51 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=24.3, 8.7 Hz), 7.66 (dd, 2H), 8.69 (d, 1H, J=7.5 Hz).

20

Example 140

4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzamide
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-cyanobenzoyl chloride (38.6 mg, 0.23 mmol), and N,N-diisopropylethylamine (50 μL, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62% yield).
 MS (APCI+) m/z 451 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.22-2.47 (m, 6H), 2.69 (m, 1H), 3.01 (m, 1H), 3.19 (m, 1H), 3.41 (m, 2H), 3.91 (m, 1H), 4.05 (m, 1H), 4.17 (t, 2H, J=5.1 Hz), 5.18 (m, 1H), 6.96 (d, 2H, J=8.5 Hz), 7.52 (d, 2H, J=8.5 Hz), 7.66 (dd, 4H, J=24.7, 8.5 Hz), 7.73 (d, 2H, J=8.8 Hz), 8.10 (d, 2H, J=8.1 Hz), 9.05 (m, 1H).

Example 141N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)nicotinamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), nicotinoyl chloride (41.5 mg, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (65% yield).
 MS (APCI+) m/z 427 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.39 (m, 3H), 2.71 (m, 1H), 3.07 (m, 1H), 3.24 (m, 1H), 3.43 (m, 2H), 3.93 (m, 1H), 4.05 (m, 1H), 4.17 (t, 2H, J=5.1 Hz), 6.96 (d, 2H, J=9.2 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.66 (dd, 4H, J=24.1, 8.5 Hz), 7.77 (dd, 1H, J=8.2, 5.4 Hz), 8.81 (dt, 1H, J=8.1, 1.7 Hz), 8.88 (dd, 1H, J=5.5, 1.4 Hz), 9.36 (d, 1H, J=1.7 Hz), 9.60 (d, 1H, J=8.2 Hz).

15

Example 142N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-furamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 2-furoyl chloride (0.023 ml, 0.22 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (68% yield).
 MS (APCI+) m/z 416 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.29-2.49 (m, 4H), 2.57-2.79 (m, 2H), 2.79-3.05 (m, 5H), 3.07-3.28 (m, 1H), 3.37 (m, 1H), 3.46-3.73 (m, 4H), 3.89 (m, 1H), 4.17 (m, 4H), 5.35 (m, 1H), 6.45 (m, 2H), 6.98 (m, 2H), 7.11 (d, 2H, J=3.4 Hz), 7.42 (m, 2H), 7.53 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=20.2, 8.3 Hz).

Example 143N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 2-(3-pyridinyl)-1,3-thiazole-4-carbonyl chloride (48.1 mg, 0.23 mmol), and N,N-

diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (72% yield).

5 MS (APCI+) m/z 510 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.4 (m, 2H), 2.65 (m, 1H), 3.0 (m, 1H), 3.2 - 3.6 (m, 2H), 4.0 (m, 1H), 4.20 (m, 2H), 4.95 (m, 4H), 5.2 (m, 1H), 6.95 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.55 (m, 1H), 7.65 (m, 4H), 7.85 (m, 1H), 8.25 (s, 1H), 8.8 (m, 1H), 9.3 (m, 1H).

10

Example 144

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-propanesulfonamide
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 2-propanesulfonyl chloride (26.2 μ L, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (70% yield).
 15 MS (APCI+) m/z 428 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.4 (d, J=7Hz, 6H), 2.4 (m, 2H), 2.6 (m, 2H), 2.9 (m, 2H), 3.1 (m, 2H), 3.4 (m, 2H), 3.95 (m, 2H), 4.15 (m, 2H), 4.4 (m, 1H), 6.95 (d, J=9Hz, 2H), 7.50 (d, J=9Hz, 2H), 7.65 (m, 4H).

20

Example 145

N'-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N-dimethylsulfamide
 4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), dimethylsulfamoyl chloride (0.025 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (65% yield).
 25 MS (APCI+) m/z 429 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.14-2.42 (m, 8H), 2.57 (m, 1H), 2.82 (s, 6H), 3.07 (m, 1H), 3.34 (m, 2H), 3.97 (m, 2H), 4.14 (t, 2H, J=5.6 Hz), 4.36 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.54 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=21.6, 8.6 Hz).

Example 146N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl)-4-(methylsulfonyl)benzenesulfonamide

5 4'-{3-[3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-(methylsulfonyl)benzenesulfonyl chloride (59.4 mg, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (75% yield).

10 MS (APCI+) m/z 540 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 2.13 (m, 2H), 2.23-2.51 (m, 8H), 2.85 (m, 1H), 3.00 (m, 1H), 3.10 (s, 3H), 3.30 (m, 2H), 3.72 (m, 1H), 3.88 (m, 1H), 4.12 (t, 2H, J=5.4 Hz), 4.37 (m, 1H), 6.95 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=8.9 Hz), 7.66 (dd, 4H, J=22.1, 8.5 Hz), 8.11 (d, 4H, J=3.1 Hz).

15

Example 147N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl)-N-(3,3-dimethylbutanoyl)-3,3-dimethylbutanamide

20 4'-{3-[3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), tert-butylacetyl chloride (0.032 mL, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (25% yield).

25 MS (APCI+) m/z 518 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.06 (s, 18H), 2.19-2.42 (m, 3H), 2.55 (s, 2H), 2.58 (s, 2H), 3.04-3.53 (m, 8H), 3.60 (m, 1H), 3.67-4.06 (m, 2H), 4.13 (t, 2H, J=5.4 Hz), 4.92 (m, 1H), 6.97 (d, 2H, J=8.9 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=20.9, 8.6 Hz).

Example 148

30 N'-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl)-N'-(dimethylaminocarbonyl)-N,N-dimethylurea

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), dimethylcarbamyl chloride (0.021 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (30% yield).
MS (APCI+) m/z 464 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.86-2.18 (m, 3H), 2.36 (m, 1H), 2.82 (s, 6H), 2.87-3.07 (m, 10H), 3.10 (s, 3H), 3.25 (s, 3H), 3.30-3.52 (m, 4H), 4.01 (t, 2H, J=5.6 Hz), 4.15 (m, 1H), 4.34 (m, 1H), 4.50 (m, 1H), 4.87 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=9.1 Hz), 7.67 (dd, 4H, J=17.8, 8.7 Hz).

10

Example 149N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(1-pyrrolidinylcarbonyl)-1-pyrrolidinecarboxamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 1-pyrrolidinecarbonyl chloride (0.031 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (25% yield).
MS (APCI+) m/z 516 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.78 (m, 2H), 1.87-2.20 (m, 5H), 2.21-2.47 (m, 1H), 3.17-3.59 (m, 7H), 3.68 (m, 2H), 3.77-4.25 (m, 6H), 4.35 (m, 1H), 4.52 (m, 1H), 4.77-5.00 (m, 1H), 6.97 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=8.4 Hz), 7.67 (dd, 4H, J=18.3, 8.5 Hz).

25

Example 150N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(4-morpholinylcarbonyl)-4-morpholinecarboxamide

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (50 mg, 0.15 mmol), 4-morpholinecarbonyl chloride (0.027 ml, 0.23 mmol), and N,N-diisopropylethylamine (50 μ L, 0.30 mmol) in 2 mL of dichloromethane were stirred at 20 °C for 18 hours. The mixture was evaporated under reduced pressure and the residue was

purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (75% yield). ¹H NMR (300 MHz, CDCl₃) δ 2.09 (m, 2H), 2.34 (m, 1H), 3.22 (m, 3H), 3.44 (m, 6H), 3.57 (m, 4H), 3.68 (m, 7H), 3.80 (m, 5H), 4.02 (t, 2H, J=5.4 Hz), 4.13 (m, 1H), 4.36 (m, 1H), 4.53 (m, 1H), 4.90 (m, 1H), 6.96 (d, 2H, J=8.5 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.67 (dd, 4H, J=19.3, 8.8 Hz). MS (APCI+) m/z 548 (M+H)⁺.

Example 151

cyclopropyl{4-[3-(3-hydroxy-1-pyrrolidinyl)propoxy]phenyl}methanone

10

Example 151A

cyclopropyl(4-hydroxyphenyl)methanone

Sodium hydroxide in water (50% (w/w), 40.4 mL) was treated over a period of 15 minutes with para-hydroxy-4-chlorobutyrophenone (25.0 g, 137 mmol), followed by additional aqueous sodium hydroxide (25% (w/w), 177 mL). Additional para-hydroxy-4-chlorobutyrophenone (25.0 g, 137 mmol) was added portionwise to the reaction mixture, followed by solid sodium hydroxide (40.4 g). A yellow precipitate formed. After refluxing for 60 minutes, water (50 mL) was added and the resulting mixture was refluxed for another 60 minutes, cooled to room temperature, diluted with water (100 mL) and neutralized with acetic acid. The precipitate was collected by filtration, washed with water, air-dried, and triturated at 40 °C with chloroform (1.5 L). The chloroform solution was dried (MgSO₄), filtered, and concentrated. The concentrate was recrystallized from chloroform/hexanes to afford 26.65 g (95%) of the desired product.

MS (APCI-) m/z 161 (M-H)⁻; ¹H NMR (300 MHz, CDCl₃) δ 7.9 (d, 2H), 6.9 (d, 2H), 2.65 (m, 1H), 1.23 (m, 2H), 1.03 (m, 2H).

25

Example 151B

(4-(3-chloropropoxy)phenyl)(cyclopropyl)methanone

The product from Example 151A (10 g, 61.7 mmol), K₂CO₃ (12.7 g, 91.9 mmol), and 1-bromo-3-chloropropane (10.74 g, 68.2 mmol) in 2-butanone (100 mL) was refluxed for 24 hours, cooled to ambient temperature, filtered, and concentrated. The filtrate was heated at 40 °C under reduced pressure for 3 hours to provide the title compound (13.256 g, 90% yield) of sufficient purity for subsequent use without further purification.

Example 151Ccyclopropyl(4-[3-(3-hydroxy-1-pyrrolidinyl)propoxy]phenyl)methanone

The product from Example 151B (200 mg, 0.83 mmol), 3-hydroxypyrrolidine (72 mg, 0.83 mmol), potassium carbonate (121 mg) and potassium iodide (146 mg) in 10 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (68 % yield).

MS (ESI+) m/z 290 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 0.93 (m, 2H), 1.20 (m, 2H), 4.18 (t, J=5 Hz, 2H), 4.42 (s, 1H), 6.90 (d, J=7 Hz, 2H), 8.00 (d, J=7 Hz, 2H).

Example 152cyclopropyl(4-{3-[(3R)-3-hydroxypyrrolidinyl]propoxy}phenyl)methanone

(4-(3-Chloropropoxy)phenyl)(cyclopropyl)methanone (150 mg, 0.63 mmol), 3-(3R)-hydroxypyrrolidine (55 mg, 0.63 mmol), potassium carbonate (152 mg) and potassium iodide (183 mg) in 25 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (62 % yield).

MS (ESI+) m/z 290 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 0.95 (m, 2H), 1.21 (m, 2H), 4.13 (t, J=5 Hz, 2H), 4.40 (s, 1H), 6.95 (d, J=7 Hz, 2H), 8.00 (d, J=7 Hz, 2H).

Example 1534-{3-[(3R)-3-hydroxypyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile

4-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (200 mg, 0.74 mmol), (3R)-3-pyrrolidinol (70 mg, 0.81 mmol), potassium carbonate (152 mg) and potassium iodide (183 mg) in 25 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (69 % yield).

MS (ESI+) m/z 323 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 3.0 (m, 1H), 4.13 (t, J=5 Hz, 2H), 4.40 (s, 1H), 7.0 (d, J=7 Hz, 2H), 7.55 (d, J=7 Hz, 2H), 7.65 (m, 4H).

Example 1544'-[3-(3-oxo-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile

Oxalyl chloride (0.195 mmol) in dry CH_2Cl_2 (5 mL) under an atmosphere of nitrogen at -78 °C was treated with DMSO (0.39 mmol). After 15 minutes of stirring, the product 5 from Example 153 was added and the mixture was stirred for another 15 minutes. Triethylamine (0.45 mmol) was added and the mixture was gradually allowed to warm to 0 °C. The solution was partitioned between water (50 mL) and dichloromethane (50 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate and filtered. The filtrate was evaporated under reduced pressure and the residue purified by chromatography 10 ($\text{CHCl}_3:\text{MeOH}:\text{NH}_4\text{OH}$, 9:1:0.1) to provide the title compound (60% yield).
MS (ESI+) m/z 321 ($\text{M}+\text{H}$)⁺; ¹H NMR (300 MHz, CDCl_3) δ 4.15 (t, $J=5$ Hz, 2H), 7.00 (d, $J=7$ Hz, 2H), 7.57 (d, $J=7$ Hz, 2H), =7.63 (m, 4H).

Example 155

15 4'-(3-[3S)-3-hydroxypyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile
4'-(3-[3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (180 mg, 0.65 mmol), (3S)-3-pyrrolidinol (57 mg, 0.65 mmol), potassium carbonate (140 mg) and potassium iodide (180 mg) in 20 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by 20 chromatography ($\text{CHCl}_3:\text{MeOH}:\text{NH}_4\text{OH}$, 9:1:0.1) to provide the title compound (61 % yield).
MS (ESI+) m/z 323 ($\text{M}+\text{H}$)⁺; ¹H NMR (300 MHz, CDCl_3) δ 2.02 (m, 2H), 2.25 (m, 3H), 4.15 (t, $J=5$ Hz, 2H), 6.99 (d, $J=7$ Hz, 2H), 7.55 (d, $J=7$ Hz, 2H), 7.65 (m, 4H).

Example 156

25 4'-(3-(3-hydroxy-3-methyl-1-pyrrolidinyl)propoxy)[1,1'-biphenyl]-4-carbonitrile
The product from Example 154 (130 mg, 0.41 mmol) in dry THF (6 mL) under an atmosphere of nitrogen at 0 °C was treated with a 3.0M solution of methylmagnesium bromide (0.82 mmol). After warming to ambient temperature and stirring for 2 hours, the mixture was quenched with 3 mL of cold saturated ammonium chloride solution and then the 30 solution was partitioned between water (50 mL) and dichloromethane (50 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate and filtered. The filtrate was

evaporated under reduced pressure and the residue purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (55% yield).

MS (ESI+) m/z 337 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.22 (s, 3H), 1.93 (m, 2H), 2.1 (m, 2H), 2.82 (m, 2H), 2.92 (d, J=8 Hz, 1H), 3.23 (m, 1H), 4.05 (t, J=5 Hz, 2H), 7.00 (d, J=7 Hz, 2H), 7.57 (d, J=7 Hz, 2H), 7.63 (m, 4H).

Example 157

4'-[3-(3-hydroxy-3-isopropyl-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile

The product from Example 154 (147 mg, 0.46 mmol) in dry THF (10 mL) under an atmosphere of nitrogen at 0 °C was treated with a 2.0M solution of isopropylmagnesium chloride (0.92 mmol). After warming to ambient temperature and stirring for 1 hour, the mixture was quenched with 5 mL of cold saturated ammonium chloride solution and then the solution was partitioned between water (50 mL) and dichloromethane (50 mL). The organic layer was washed with brine (50 mL), dried over sodium sulfate and filtered. The filtrate was evaporated under reduced pressure and the residue purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (75% yield).

MS (ESI+) m/z 365 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 4.25 (t, J=5 Hz, 2H), 6.98 (d, J=7 Hz, 2H), 7.55 (d, J=7 Hz, 2H), 7.62 (m, 4H).

20

Example 158

4'-{3-[(3R)-3-hydroxy-3-methylpyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile

4'-{3-[(3R)-3-Aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile (177 mg, 0.65 mmol), (3R)-3-methyl-3-pyrrolidinol (75 mg, 0.74 mmol), potassium carbonate (153 mg) and potassium iodide (184 mg) in 10 mL of DMF were heated at 110 °C for 10 hours.

25 The mixture was evaporated under reduced pressure and the residue was purified by chromatography (CHCl₃:MeOH:NH₄OH, 9:1:0.1) to provide the title compound (40 % yield). MS (ESI+) m/z 337 (M+H)⁺; ¹H NMR (300 MHz, CDCl₃) δ 1.2 (s, 3H), 4.15 (t, J=5 Hz, 2H), 6.98 (d, J= 7 Hz, 2H), 7.55 (d, J=7 Hz, 2H), 7.69 (m, 4H).

30

Example 159

N,N-dimethyl-N-[(3R)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl]amine

1-{{4'-(3-Chloropropoxy)[1,1'-biphenyl]-4-yl]carbonyl}pyrrolidine (200 mg, 0.74 mmol), N,N-dimethyl-N-[(3R)-pyrrolidinyl]amine (70 mg, 0.81 mmol), potassium carbonate (152 mg) and potassium iodide (183 mg) in 25 mL of 2-butanone were heated at 110 °C for 72 hours. The mixture was evaporated under reduced pressure and the residue was purified by HPLC chromatography (0 to 95% CH₃CN in H₂O with 0.1% TFA, 10 minute linear gradient) to provide the title compound in 79% yield.

10 ¹HNMR (500 MHz, CDCl₃) δ 1.89 (m, 2H), 1.97 (m, 2H), 2.26 (m, 2H), 2.57 (m, 3H), 2.86 (s, 6H), 3.40 (m, 3H), 3.49 (t, 2H, J=6.7 Hz), 3.67 (t, 2H, J=6.8 Hz), 3.74 (m, 1H), 3.84 (dd, 1H, J=10.6, 9.0 Hz), 4.02 (m, 1H), 4.10 (t, 2H, J=5.6 Hz), 4.17 (m, 1H), 6.95 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=8.8 Hz), 7.57 (s, 4H); MS (APCI) m/z 422 (M+H)⁺.

Example 160

15 N,N-dimethyl-N-[(3S)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl]amine

1-{{4'-(3-Chloropropoxy)[1,1'-biphenyl]-4-yl]carbonyl}pyrrolidine and N,N-dimethyl-N-[(3S)-pyrrolidinyl]amine were processed as described in Example 159 to provide the title compound.

20 ¹HNMR (500 MHz, CDCl₃) δ 1.88 (m, 3H), 1.98 (m, 2H), 2.26 (m, 3H), 2.56 (m, 3H), 2.86 (s, 6H), 3.45 (m, 5H), 3.70 (m, 3H), 3.84 (m, 1H), 4.03 (m, 1H), 4.10 (t, 2H, J=5.6 Hz), 4.19 (m, 1H), 6.95 (d, 2H, J=8.8 Hz), 7.53 (d, 2H, J=9.1 Hz), 7.57 (s, 4H); MS (APCI) m/z 422 (M+H)⁺.

Example 161

(3R)-1-(3-{[4'-(1-pyrrolidinylcarbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)-3-pyrrolidinol

1-{{4'-(3-Chloropropoxy)[1,1'-biphenyl]-4-yl]carbonyl}pyrrolidine and (3R)-3-pyrrolidinol were processed as described in Example 159 to provide the title compound.

30 ¹HNMR (500 MHz, CDCl₃) δ 1.82-2.02 (m, 5H), 2.03-2.50 (m, 16H), 2.96 (m, 2H), 3.14-3.44 (m, 5H), 3.49 (t, 2H, J=6.5 Hz), 3.59 (m, 1H), 3.67 (t, 2H, J=6.9 Hz), 3.87 (m, 1H), 3.98

(m, 1H), 4.06 (m, 1H), 4.12 (m, 1H), 4.64 (m, 1H), 6.94 (d, 2H, J=8.7 Hz), 7.53 (d, 2H, J=8.7 Hz), 7.57 (s, 4H); MS (APCI) m/z 395 (M+H)⁺.

Example 162

5 N,N-dimethyl-N-[(3R)-1-(3-{4-[4-(1-pyrrolidinylcarbonyl)-1-piperazinyl]phenoxy}propyl)pyrrolidinyl]amine

1-[4-(3-chloropropoxy)phenyl]-4-(1-pyrrolidinylcarbonyl)piperazine and N,N-dimethyl-N-[(3R)-pyrrolidinyl]amine were processed as described in Example 159 to provide the title compound.

10 ¹HNMR (500 MHz, CDCl₃) δ 1.86 (m, 4H), 2.23 (m, 4H), 2.58 (m, 4H), 2.86 (s, 6H), 3.25 (m, 4H), 3.43 (m, 6H), 3.61 (m, 4H), 3.72 (m, 1H), 3.85 (m, 1H), 4.05 (m, 3H), 4.18 (m, 1H), 6.88 (d, 2H, J=9.0 Hz), 7.17 (d, 2H, J=9.0 Hz); MS (APCI) m/z 430 (M+H)⁺.

Example 163

15 N,N-dimethyl-N-[(3S)-1-(3-{4-[4-(1-pyrrolidinylcarbonyl)-1-piperazinyl]phenoxy}propyl)pyrrolidinyl]amine

1-[4-(3-chloropropoxy)phenyl]-4-(1-pyrrolidinylcarbonyl)piperazine and N,N-dimethyl-N-[(3S)-pyrrolidinyl]amine were processed as described in Example 159 to provide the title compound.

20 ¹HNMR (500 MHz, CDCl₃) δ 1.86 (m, 4H), 2.22 (m, 4H), 2.57 (m, 4H), 2.86 (s, 6H), 3.25 (m, 4H), 3.42 (m, 6H), 3.61 (m, 4H), 3.73 (m, 1H), 3.85 (m, 1H), 4.04 (m, 3H), 4.18 (m, 1H), 6.88 (d, 2H, J=9.0 Hz), 7.17 (d, 2H, J=9.0 Hz); MS (APCI) m/z 430 (M+H)⁺.

Determination of Biological Activity

25 Histamine-3 Receptor Binding

To determine the effectiveness of representative compounds of this invention as histamine-3 receptor ligands (H₃ receptor ligands), the following tests were conducted according to methods previously described (European Journal of Pharmacology, 188:219-227 (1990); Journal of Pharmacology and Experimental Therapeutics, 275: 598-604 (1995); Journal of Pharmacology and Experimental Therapeutics, 276:1009-1015 (1996); and Biochemical Pharmacology, 22: 3099-3108 (1973)).

100

Briefly, male Sprague-Dawley rat brain cortices were homogenized (1 g tissue/10 mL buffer) in 50 mM Tris-HCl/5 mM EDTA containing protease inhibitor cocktail (Calbiochem) using a polytron set at 20,500 rpm. Homogenates were centrifuged for 20 minutes at 40,000 \times g. The supernatant was decanted, and pellets were weighed. The pellet was 5 resuspended by polytron homogenization in 40 mL 50 mM Tris-HCl/5 mM EDTA with protease inhibitors and centrifuged for 20 minutes at 40,000 \times g. The membrane pellet was resuspended in 6.25 volumes (per gram wet weight of pellet) of 50 mM Tris-HCl/5 mM EDTA with protease inhibitors and aliquots flash frozen in liquid N₂ and stored at -70 °C until used in assays. Rat cortical membranes (12 mg wet weight/tube) were incubated with (³H)- 10 N- α -methylhistamine (~0.6 nM) with or without H₃ receptor antagonists in a total incubation volume of 0.5 mL of 50 mM Tris-HCl/5 mM EDTA (pH 7.7). Test compounds were dissolved in DMSO to provide a 20 mM solution, serially diluted and then added to the incubation mixtures prior to initiating the incubation assay by addition of the membranes. 15 Thioperamide (3 μ M) was used to determine nonspecific binding. Binding incubations were conducted for 30 minutes at 25 °C and terminated by addition of 2 mL of ice cold 50 mM Tris-HCl (pH 7.7) and filtration through 0.3% polyethylenimine-soaked Unifilter plates (Packard). These filters were washed 4 additional times with 2 mL of ice-cold 50 mM Tris-HCl and dried for 1 hour. Radioactivity was determined using liquid scintillation counting 20 techniques. Results were analyzed by Hill transformation and Ki values were determined using the Cheng-Prusoff equation.

Table 1

Example Number	Ki (nM)
1	105
2	68
3	37
4	55
5	6.3
6	95
7	31
8	1721

9	675
10	626
11	4.0
12	71
13	50
14	40
15	7.3
16	67
17	13
18	29
19	3.8
20	12
21	22
22	12
23	33
24	43
25	30
26	57
27	18
28	16
29	5.9
30	10
31	30
32	99
33	16
34	11
35	13
36	12
37	349
38	132

39	7.6
40	51
41	48
42	8.8
43	37
44	52
45	127
46	103
47	12
48	60
49	13
50	60
51	9.3
52	15
53	3.9
54	98
55	2.9
56	4.7
57	3.6
58	3.9
59	3.0
60	5.2
61	4.2
62	6.4
63	12
64	11
65	5.0
66	4.6
67	2.9
68	2.8

69	3.7
70	3.0
71	2.8
72	7.9
73	6.4
74	5.6
75	19
76	4.9
77	5.3
78	8.7
79	7.9
80	10
81	5.0
82	21
83	23
84	9.5
85	5.3
86	1.8
87	4.9
88	6.8
89	11
90	11
91	2.8
92	4.9
93	3.2
94	4.9
95	4.6
96	1.2
97	6.7
98	90

99	63
100	1634
101	179
102	112
103	195
104	173
105	725
106	168
107	200
108	12
109	26
110	30
111	11
112	13
113	13
114	65
115	21
116	14
117	45
118	5.7
119	7.7
120	10
121	16
122	20
123	5.6
124	2.9
125	7.4
126	4.6
127	16
128	2.3

129	192
130	31
131	51
132	74
133	57
134	168
135	132
136	215
137	196
138	118
139	288
140	220
141	342
142	321
143	85
144	21
145	9.6
146	3.0
147	209
148	291
149	329
150	267
151	22
152	8.7
153	42
154	63
155	105
156	45
157	593
158	15

As shown by the data in Table 1, the compounds of the present invention bind to the histamine-3 receptor and therefore may have utility in the treatment of diseases or conditions ameliorated with histamine-3 receptor ligands.

Compounds of the present invention may exist as stereoisomers wherein, asymmetric or chiral centers are present. These stereoisomers are "R" or "S" depending on the configuration of substituents around the chiral carbon atom. The terms "R" and "S" used herein are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem., 1976, 45: 13-30. In particular, the stereochemistry at the point of attachment of R₆ and R₉, as shown in formula I-VII, may 5 independently be either (R) or (S), unless specifically noted otherwise. Additionally, the 3-position of the pyrrolidine ring of formula I-VII may independently be either (R) or (S), unless specifically noted otherwise. The preferred stereochemistry at the 3-position of the pyrrolidine ring is the (R) configuration for compounds of the present invention. However, the present invention does contemplate various stereoisomers and mixtures thereof and are 10 specifically included within the scope of this invention. Stereoisomers include enantiomers and diastereomers, and mixtures of enantiomers or diastereomers. Individual stereoisomers of compounds of the present invention may be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of 15 racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct 20 separation of the mixture of optical enantiomers on chiral chromatographic columns.

The term "pharmaceutically acceptable carrier," as used herein, means a non-toxic, 25 inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as 30 cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols; such a propylene glycol; esters such as

ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, 5 releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of one skilled in the art of formulations.

10 The present invention provides pharmaceutical compositions which comprise compounds of the present invention formulated together with one or more non-toxic pharmaceutically acceptable carriers. The pharmaceutical compositions can be formulated for oral administration in solid or liquid form, for parenteral injection or for rectal administration.

15 Further included within the scope of the present invention are pharmaceutical compositions comprising one or more of the compounds of formula I-VII prepared and formulated in combination with one or more non-toxic pharmaceutically acceptable compositions. The pharmaceutical compositions can be formulated for oral administration in solid or liquid form, for parenteral injection or for rectal administration.

20 The pharmaceutical compositions of this invention can be administered to humans and other mammals orally, rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments or drops), buccally or as an oral or nasal spray. The term "parenterally," as used herein, refers to modes of administration which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous, intraarticular injection and infusion.

25 Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions and sterile powders for reconstitution into sterile injectable solutions or dispersions. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (propylene glycol, polyethylene glycol, glycerol, and the like), suitable mixtures thereof, vegetable oils (such as olive oil) and 30 injectable organic esters such as ethyl oleate. Proper fluidity may be maintained, for

example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersions, and by the use of surfactants.

These compositions may also contain adjuvants such as preservative agents, wetting agents, emulsifying agents, and dispersing agents. Prevention of the action of 5 microorganisms may be ensured by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, and the like. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form may be brought about by the use of agents delaying absorption, for example, aluminum monostearate and gelatin.

10 In some cases, in order to prolong the effect of a drug, it is often desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution which, in turn, may depend upon crystal size and crystalline form. Alternatively, 15 delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

20 Suspensions, in addition to the active compounds, may contain suspending agents, as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, tragacanth, and mixtures thereof.

If desired, and for more effective distribution, the compounds of the present invention can be incorporated into slow-release or targeted-delivery systems such as polymer matrices, 25 liposomes, and microspheres. They may be sterilized, for example, by filtration through a bacteria-retaining filter or by incorporation of sterilizing agents in the form of sterile solid compositions, which may be dissolved in sterile water or some other sterile injectable medium immediately before use.

30 The active compounds can also be in micro-encapsulated form, if appropriate, with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound can be admixed with at least one inert diluent

such as sucrose, lactose, or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may 5 optionally contain opacifying agents and can also be of such composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

Injectable depot forms are made by forming microencapsulated matrices of the drug 10 in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions which are compatible with body tissues.

15 The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

20 Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic, parenterally acceptable diluent or solvent such as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution.

25 In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

30 Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium

phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and salicylic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidinone, sucrose, and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay; and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract in a delayed manner. Examples of embedding compositions which can be used include polymeric substances and waxes.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol,

tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof.

Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, ear drops, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

The ointments, pastes, creams and gels may contain, in addition to an active compound of this invention, excipients such as animal and vegetable fats, oils, waxes, paraffins, starch, tragacanth, cellulose derivatives, polyethylene glycols, silicones, bentonites, silicic acid, talc and zinc oxide, or mixtures thereof.

Powders and sprays can contain, in addition to the compounds of this invention, excipients such as lactose, talc, silicic acid, aluminum hydroxide, calcium silicates and polyamide powder, or mixtures of these substances. Sprays can additionally contain customary propellants such as chlorofluorohydrocarbons.

Compounds of the present invention may also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes may be used. The present compositions in liposome form may contain, in addition to the compounds of the present invention, stabilizers, preservatives, excipients, and the like. The preferred lipids are the natural and synthetic phospholipids and phosphatidylcholines (lecithins) used separately or together.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., Methods in Cell Biology, Volume XIV, Academic Press, New York, N. Y., (1976), p 33 et seq.

The terms "pharmaceutically acceptable salts, esters and amides," as used herein, refer to carboxylate salts, amino acid addition salts, zwitterions, esters and amides of compounds of formula I-VII which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue 5 toxicity, irritation, allergic response, and the like, are commensurate with a reasonable benefit/risk ratio, and are effective for their intended use.

The compounds of the present invention can be used in the form of pharmaceutically acceptable salts derived from inorganic or organic acids. By "pharmaceutically acceptable salt" is meant those salts which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, 10 irritation, allergic response and the like and are commensurate with a reasonable benefit/risk ratio. Pharmaceutically acceptable salts are well-known in the art. For example, S. M. Berge et al. describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 1977, 66: 1 et seq. The salts can be prepared in situ during the final isolation and purification of the 15 compounds of the invention or separately by reacting a free base function with a suitable organic acid. Representative acid addition salts include, but are not limited to acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethansulfonate 20 (isethionate), lactate, maleate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate and undecanoate. Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl 25 halides such as methyl, ethyl, propyl, and butyl chlorides, bromides and iodides; dialkyl sulfates like dimethyl, diethyl, dibutyl and diethyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; arylalkyl halides like benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained. Examples of acids which can be employed to form pharmaceutically acceptable 30 acid addition salts include such inorganic acids as hydrochloric acid, hydrobromic acid, sulphuric acid and phosphoric acid and such organic acids as oxalic acid, maleic acid, succinic acid and citric acid.

Basic addition salts can be prepared in situ during the final isolation and purification of compounds of this invention by reacting a carboxylic acid-containing moiety with a suitable base such as the hydroxide, carbonate or bicarbonate of a pharmaceutically acceptable metal cation or with ammonia or an organic primary, secondary or tertiary amine.

5 Pharmaceutically acceptable salts include, but are not limited to, cations based on alkali metals or alkaline earth metals such as lithium, sodium, potassium, calcium, magnesium and aluminum salts and the like and nontoxic quaternary ammonia and amine cations including ammonium, tetramethylammonium, tetraethylammonium, methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine and the like. Other representative 10 organic amines useful for the formation of base addition salts include ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine and the like. Preferred salts of the compounds of the invention include phosphate, tris and acetate.

The term "pharmaceutically acceptable ester," as used herein, refers to esters of compounds of the present invention which hydrolyze in vivo and include those that break 15 down readily in the human body to leave the parent compound or a salt thereof. Examples of pharmaceutically acceptable, non-toxic esters of the present invention include C₁-to-C₆ alkyl esters and C₅-to-C₇ cycloalkyl esters, although C₁-to-C₄ alkyl esters are preferred. Esters of the compounds of formula I-VII may be prepared according to conventional methods.

The term "pharmaceutically acceptable amide," as used herein, refers to non-toxic 20 amides of the present invention derived from ammonia, primary C₁-to-C₆ alkyl amines and secondary C₁-to-C₆ dialkyl amines. In the case of secondary amines, the amine may also be in the form of a 5- or 6-membered heterocycle containing one nitrogen atom. Amides derived from ammonia, C₁-to-C₃ alkyl primary amides and C₁-to-C₂ dialkyl secondary 25 amides are preferred. Amides of the compounds of formula I-VII may be prepared according to conventional methods.

The term "pharmaceutically acceptable prodrug" or "prodrug," as used herein, represents those prodrugs of the compounds of the present invention which are, within the 30 scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use. Prodrugs of the present invention may be rapidly transformed in vivo to a parent compound

of formula I-VII, for example, by hydrolysis in blood. A thorough discussion is provided in T. Higuchi and V. Stella, Pro-drugs as Novel Delivery Systems, V. 14 of the A.C.S. Symposium Series, and in Edward B. Roche, ed., Bioreversible Carriers in Drug Design, American Pharmaceutical Association and Pergamon Press (1987), hereby incorporated by reference. Representative examples of prodrugs of the present invention include, but are not limited to, (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl methyl sulfate, (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl diethyl phosphate, (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl acetate, (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl dimethylcarbamate, and (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl methyl carbonate.

Dosage forms for topical administration of a compound of this invention include powders, sprays, ointments and inhalants. The active compound is mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers or propellants which can be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention can be varied so as to obtain an amount of the active compound(s) which is effective to achieve the desired therapeutic response for a particular patient, compositions and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity of the condition being treated and the condition and prior medical history of the patient being treated. However, it is within the skill of the art to start doses of the compound at levels lower than required for to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The present invention contemplates pharmaceutically active compounds either chemically synthesized or formed by in vivo biotransformation to compounds of formula I-VII.

The compounds of the present invention, including but not limited to those specified in the examples, possess an affinity for the histamine-3 receptor. As histamine-3 receptor ligands, the compounds of the present invention may be useful for the treatment and prevention of diseases or conditions such as acute myocardial infarction, Alzheimer's disease,

attention-deficit hyperactivity disorder, Parkinson's disease, epilepsy, schizophrenia, depression, cutaneous carcinoma, medullary thyroid carcinoma, melanoma, asthma, narcolepsy, Meniere's disease, gastrointestinal disorders, inflammation, migraine, motion sickness, obesity, pain, seizures, and septic shock.

5 The ability of the compounds of the present invention, including but not limited to those specified in the examples, to treat septic shock and cardiovascular disorders, in particular, acute myocardial infarction may be demonstrated by (Imamura et al., Circ.Res., (1996) 78, 475-481; Imamura et al., Circ.Res., (1996) 78, 863-869; R. Levi and N.C.E. Smith, "Histamine H₃-receptors: A new frontier in myocardial ischemia", J. Pharm. Exp. Ther., 292: 825-830, (2000); and Hatta, E., K. Yasuda and R. Levi, "Activation of histamine H₃ receptors inhibits carrier-mediated norepinephrine release in a human model of protracted myocardial ischemia", J. Pharm. Exp. Ther., 283: 494-500, (1997)).

10

15 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat sleep disorders, in particular, narcolepsy may be demonstrated by (Lin et al., Brain Res. (1990) 523, 325-330; Monti et al., Neuropsychopharmacology (1996) 15, 31-35; Sakai, et al., Life Sci. (1991) 48, 2397-2404; Mazurkiewicz-Kwilecki and Nsonwah, Can. J. Physiol. Pharmacol. (1989) 67, 75-78; Panula, P. et al., Neuroscience (1998) 44, 465-481); Wada et al., Trends in Neuroscience (1991) 14, 415); and Monti et al., Eur. J. Pharmacol. (1991) 205, 283).

20 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat cognition and memory process disorders may be demonstrated by (Mazurkiewicz-Kwilecki and Nsonwah, Can. J. Physiol. Pharmacol. (1989) 67, 75-78; Panula, P. et al., Neuroscience (1998) 44, 465-481; Haas et al., Behav. Brain Res. (1995) 66, 41-44; De Almeida and Izquierdo, Arch. Int. Pharmacodyn. (1986) 283, 193-198; Kamei et al., Psychopharmacology (1990) 102, 312-318; and Kamei and Sakata, Jpn. J. Pharmacol. (1991) 57, 437-482); Schwartz et al., Psychopharmacology; The fourth Generation of Progress. Bloom and Kupfer (eds). Raven Press, New York, (1995) 397; and Wada et al., Trends in Neurosci., (1991) 14, 415).

25

30 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat attention-deficit hyperactivity disorder (ADHD) may be demonstrated by (Shaywitz et al., Psychopharmacology (1984) 82, 73-77; Dumery and

Blozovski, Exp. Brain Res. (1987) 67, 61-69; Tedford et al., J. Pharmacol. Exp. Ther. (1995) 275, 598-604; and Tedford et al., Soc. Neurosci. Abstr. (1996) 22, 22).

The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat seizures, in particular, epilepsy may be demonstrated by

5 (Yokoyama et al., Eur. J. Pharmacol. (1993) 234, 129; Yokoyama and Iinuma, CNS Drugs (1996) 5, 321; Onodera et al., Prog. Neurobiol. (1994) 42, 685; R. Leurs, R.C. Vollinga and H. Timmerman, "The medicinal chemistry and therapeutic potentials of ligand of the histamine H₃ receptor", Progress in Drug Research 45: 170-165, (1995); Leurs and Timmerman, Prog. Drug Res. (1992) 39, 127; The Histamine H₃ Receptor, Leurs and

10 Timmerman (eds), Elsevier Science, Amsterdam, The Netherlands (1998); H. Yokoyama and K. Iinuma, "Histamine and Seizures: Implications for the treatment of epilepsy", CNS Drugs, 5(5); 321-330, (1995); and K. Hurukami, H. Yokoyama, K. Onodera, K. Iinuma and T. Watanabe, AQ-0145, "A newly developed histamine H₃ antagonist, decreased seizure

15 susceptibility of electrically induced convulsions in mice", Meth. Find. Exp. Clin. Pharmacol., 17(C): 70-73, (1995)).

The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat motion sickness, Alzheimer's disease, and Parkinson's disease may be demonstrated by (Onodera et al., Prog. Neurobiol. (1994) 42, 685; Leurs and Timmerman, Prog. Drug Res. (1992) 39, 127; and The Histamine H₃ Receptor, Leurs and

20 Timmerman (eds), Elsevier Science, Amsterdam, The Netherlands (1998)).

The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat narcolepsy, schizophrenia, depression, and dementia may be demonstrated by (R. Leurs, R.C. Vollinga and H. Timmerman, "The medicinal chemistry and therapeutic potentials of ligand of the histamine H₃ receptor", Progress in Drug Research 45: 170-165, (1995); and The Histamine H₃ Receptor, Leurs and Timmerman (eds), Elsevier Science, Amsterdam, The Netherlands (1998)).

The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat obesity may be demonstrated by (Leurs et al., Trends in Pharm. Sci. (1998) 19, 177-183).

The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat inflammation and pain may be demonstrated by (Phillips et al., Annual Reports in Medicinal Chemistry (1998) 33, 31-40).

5 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat migraine may be demonstrated by (R. Leurs, R.C. Vollinga and H. Timmerman, "The medicinal chemistry and therapeutic potentials of ligand of the histamine H₃ receptor", Progress in Drug Research 45: 170-165, (1995); and Matsubara et al., Eur. J. Pharmacol. (1992) 224, 145; and Rouleau et al., J. Pharmacol. Exp. Ther. (1997) 281, 1085).

10 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat cancer, in particular, melanoma, cutaneous carcinoma and medullary thyroid carcinoma may be demonstrated by (Polish Med. Sci. Mon., (1998) vol. 4, issue 5, 747; Adam Szelag, "Role of histamine H₃-receptors in the proliferation of neoplastic cells in vitro", Med. Sci. Monit., 4(5): 747-755, (1998); and Fitzsimons, C., H. Duran, F. 15 Labombarda, B. Molinari and E. Rivera, "Histamine receptors signalling in epidermal tumor cell lines with H-ras gene alterations", Inflammation Res., 47 (Suppl 1): S50-S51, (1998)).

20 The ability of the compounds of the invention, including but not limited to those specified in the examples, to treat vestibular dysfunctions, in particular, Meniere's disease may be demonstrated by (R. Leurs, R.C. Vollinga and H. Timmerman, "The medicinal chemistry and therapeutic potentials of ligand of the histamine H₃ receptor", Progress in Drug Research 45: 170-165, (1995)).

25 The ability of the compounds of the present invention, including but not limited to those specified in the examples, to treat asthma may be demonstrated by (Delaunois A., Gustin P., Garbarg M., and Ansay M., "Modulation of acetylcholine, capsaicin and substance P effects by histamine H₃ receptors in isolated perfused rabbit lungs", European Journal of Pharmacology 277(2-3):243-50, (1995); and (Dimitriadou, et al., "Functional relationship between mast cells and C-sensitive nerve fibres evidenced by histamine H₃-receptor modulation in rat lung and spleen", Clinical Science. 87(2):151-63, (1994)).

30 Aqueous liquid compositions of the present invention are particularly useful for the treatment and prevention of asthma, epilepsy, Raynaud's syndrome, male sexual dysfunction,


female sexual dysfunction, migraine, pain, eating disorders, urinary incontinence, functional bowel disorders, neurodegeneration and stroke.

When used in the above or other treatments, a therapeutically effective amount of one of the compounds of the present invention can be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt, ester, amide or prodrug form. Alternatively, the compound can be administered as a pharmaceutical composition containing the compound of interest in combination with one or more pharmaceutically acceptable excipients. The phrase "therapeutically effective amount" of the compound of the invention means a sufficient amount of the compound to treat disorders, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgement. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts. For example, it is well within the skill of the art to start doses of the compound at levels lower than required to achieve the desired therapeutic effect and to gradually increase the dosage until the desired effect is achieved.

The total daily dose of the compounds of this invention administered to a human or lower animal may range from about 0.003 to about 10 mg/kg/day. For purposes of oral administration, more preferable doses can be in the range of from about 0.01 to about 5 mg/kg/day. If desired, the effective daily dose can be divided into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

WHAT IS CLAIMED IS:

1. A compound of formula I




I,

5 or a pharmaceutically acceptable salt thereof, wherein

Z is selected from the group consisting of a covalent bond and CH_2 ;

R₁ is selected from the group consisting of OR₂, NR₃R₄ and



10 R₂ is selected from the group consisting of hydrogen, alkoxy carbonyl, alkyl, alkyl carbonyl, aminocarbonyl, sulfono and phosphono;

R_3 and R_4 are independently selected from the group consisting of hydrogen, alkenyl, alkenylcarbonyl, alkenyloxycarbonyl, alkenylsulfonyl, alkoxy carbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, alkynyl, alkynylcarbonyl, alkynyloxycarbonyl, alkynylsulfonyl, aminocarbonyl, aminosulfonyl, arylalkyl, arylalkenylcarbonyl, arylalkenylsulfonyl, 15 arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylheterocyclecacbonyl, arylheterocylesulfonyl, aryloxyarylcacbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkylcarbonyl, cycloalkylalkylsulfonyl, cycloalkylcarbonyl, cycloalkylsulfonyl, formyl, heterocycle, heterocyclealkyl, heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcacbonyl, 20 heterocyclearylsulfonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcacbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl;

R_5 and R_6 are independently selected from the group consisting of hydrogen and alkyl;

25 R₇ is selected from the group consisting of hydrogen and alkyl; or

R_1 and R_7 together form ($=O$);

R_8 is selected from the group consisting of alkylcarbonyl, aryl, arylcarbonyl, arylcarbonylaryl, arylcarbonylheterocycle, cycloalkylcarbonyl, cycloalkylcarbonylaryl, cycloalkylcarbonylheterocycle, heterocycle, heterocyclecarbonyl, heterocyclecarbonylaryl, heterocyclecarbonylheterocycle;

5 R_9 is selected from the group consisting of hydrogen and lower alkyl; wherein at each occurrence of said aryl, arylalkenylcarbonyl, arylalkenylsulfonyl, arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylcarbonylaryl, arylcarbonylheterocycle, arylheterocyclecarbonyl, arylheterocylesulfonyl, aryloxyarylcarbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkylcarbonylaryl, heterocyclearylcarbonyl, heterocyclearylsulfonyl, heterocyclecarbonylaryl, heterocycleoxyarylcarbonyl, and heterocycleoxyarylsulfonyl, the aryl portion can be optionally substituted with 1, 2, or 3 substituents selected from the group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, aminoalkyl, 15 aminocarbonyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, halogen, haloalkoxy, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and nitro; and wherein at each occurrence of said arylcarbonylheterocycle, arylheterocyclecarbonyl, arylheterocylesulfonyl, cycloalkylcarbonylheterocycle, heterocycle, heterocyclealkyl, heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcarbonyl, 20 heterocyclearylsulfonyl, heterocyclecarbonyl, heterocyclecarbonylaryl, heterocyclecarbonylheterocycle, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcarbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl, the heterocycle portion can be optionally substituted with 1, 2, or 3 substituents selected from the 25 group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkyl, alkoxycarbonyl, alkyl, alkylcarbonyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfonyl, alkylthio, alkynyl, amino, aminoalkyl, aminocarbonyl, aminosulfonyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, halogen, haloalkyl, hydroxy, hydroxyalkyl, mercapto, and nitro.

30 2. A compound according to claim 1 wherein

R_1 is NR_3R_4 ;

R_3 and R_4 are independently selected from the group consisting of hydrogen, alkenyl, alkenylcarbonyl, alkenyloxycarbonyl, alkenylsulfonyl, alkoxy carbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, alkynyl, alkynylcarbonyl, alkynyloxycarbonyl, alkynylsulfonyl, aminocarbonyl, aminosulfonyl, arylalkyl, arylalkenylcarbonyl, arylalkenylsulfonyl, 5 arylalkylcarbonyl, arylalkylsulfonyl, arylarylcarbonyl, arylarylsulfonyl, arylcarbonyl, arylheterocyclecacbonyl, arylheterocylesulfonyl, aryloxyarylcacbonyl, aryloxyarylsulfonyl, arylsulfonyl, cycloalkyl, cycloalkylalkyl, cycloalkylalkylcarbonyl, cycloalkylalkylsulfonyl, cycloalkylcarbonyl, cycloalkylsulfonyl, heterocycle, heterocyclealkyl, heterocyclealkylcarbonyl, heterocyclealkylsulfonyl, heterocyclearylcacbonyl, 10 heterocyclearylsulfonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyalkylcarbonyl, heterocycleoxyarylcacbonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl; 15 R_7 is hydrogen; and
 R_8 is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

3. A compound according to claim 1 wherein

R_1 is NR_3R_4 ;
 R_3 and R_4 are independently selected from the group consisting of hydrogen, 20 alkenyloxycarbonyl, alkoxy carbonyl, alkyl, alkylcarbonyl, alkylsulfonyl, aminocarbonyl, aminosulfonyl, arylalkenylsulfonyl, arylcarbonyl, arylsulfonyl, cycloalkylcarbonyl, heterocyclecarbonyl, heterocycleheterocyclecarbonyl, heterocycleheterocyclesulfonyl, heterocycleoxyarylsulfonyl, heterocyclesulfonyl, and heterocyclethioalkylcarbonyl; 25 R_7 is hydrogen; and
 R_8 is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

4. A compound according to claim 1 wherein

Z is CH_2 ;
30 R_1 is NR_3R_4 ;

R₃ and R₄ are independently selected from the group consisting of hydrogen, alkoxy carbonyl and alkyl;

R₇ is hydrogen;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropyl carbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinyl carbonyl)phenyl, 4-(1-pyrrolidinyl carbonyl)-1-piperazinyl, 4-(1-piperidinyl carbonyl)phenyl, 4-(1-piperidinyl carbonyl)-1-piperazinyl, 4-(4-morpholinyl carbonyl)phenyl, 4-(4-morpholinyl carbonyl)-1-piperazinyl, 4-(1-azetidinyl carbonyl)phenyl, 4-(1-azetidinyl carbonyl)-1-piperazinyl, 4-(1-piperazinyl carbonyl)phenyl, 4-(1-piperazinyl carbonyl)-1-piperazinyl; and

10 R₉ is hydrogen.

5. A compound according to claim 4 selected from the group consisting of tert-butyl 1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}-3-pyrrolidinyl carbamate; N,N-dimethyl-N-[(3S)-1-(3-{[4'-(1-pyrrolidinyl carbonyl)[1,1'-biphenyl]-4-yl]oxy}propyl)pyrrolidinyl]amine; and N,N-dimethyl-N-[(3S)-1-(3-{4-[4-(1-pyrrolidinyl carbonyl)-1-piperazinyl]phenoxy}propyl)pyrrolidinyl]amine.

6. A compound according to claim 1 wherein

20 Z is a covalent bond;

R₁ is NR₂R₄;

R₃ and R₄ are independently selected from the group consisting of hydrogen, alkoxy carbonyl and alkyl;


R₇ is hydrogen;

25 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropyl carbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinyl carbonyl)phenyl, 4-(1-pyrrolidinyl carbonyl)-1-piperazinyl, 4-(1-piperidinyl carbonyl)phenyl, 4-(1-piperidinyl carbonyl)-1-piperazinyl, 4-(4-morpholinyl carbonyl)phenyl, 4-(4-morpholinyl carbonyl)-1-piperazinyl, 4-(1-azetidinyl carbonyl)phenyl, 4-(1-azetidinyl carbonyl)-1-piperazinyl, 4-(1-piperazinyl carbonyl)phenyl, 4-(1-piperazinyl carbonyl)-1-piperazinyl; and

30 R₉ is hydrogen.

7. A compound according to claim 6 that is tert-butyl (3S)-1-{2-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]ethyl}pyrrolidinylcarbamate.

5 8. A compound according to claim 1 of formula II



II,

or a pharmaceutically acceptable salt thereof wherein

10 R₁₀ is selected from the group consisting of alkenyl, alkyl, alkynyl, amino, aryl, arylalkenyl, arylalkyl, arylaryl, arylheterocycle, aryloxyaryl, cycloalkyl, cycloalkylalkyl, heterocycle, heterocyclealkyl, heteroclearyl, heterocycleheterocycle and heterocycleoxyaryl.

9. A compound according to claim 8 wherein

15 R₃ is selected from the group consisting of hydrogen and alkyl;
 R₈ is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle;
 R₁₀ is selected from the group consisting of alkyl, amino, aryl, arylalkenyl, heterocycle, heterocycleheterocycle and heterocycleoxyaryl.

20

10. A compound according to claim 8 wherein

Z is CH₂;
 R₃ is selected from the group consisting of hydrogen and alkyl;
 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-

azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

R₉ is hydrogen; and

R₁₀ is selected from the group consisting of alkyl and amino.

5

11. A compound according to claim 10 selected from the group consisting of
N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-2-propanesulfonamide;
N-{(3R)-1-[3-[4-(cyclopropylcarbonyl)phenoxy]propyl]pyrrolidinyl}-2-propanesulfonamide;

10 N-{(3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl]pyrrolidinyl}-N,N',N'-trimethyl-sulfamide;

N-{(3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-2-propanesulfonamide; and

15 N'-{(3R)-1-[3-[4'-cyano[1,1'-biphenyl]-4-yl]oxy]propyl}pyrrolidinyl)-N,N-dimethylsulfamide.

12. A compound according to claim 8 wherein

Z is CH₂;

R₃ is selected from the group consisting of hydrogen and alkyl;

20 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

25 R₉ is hydrogen; and

R₁₀ is aryl wherein said aryl is phenyl optionally substituted with 1 or 2 substituents selected from the group consisting of alkoxy, alkyl, alkylsulfonyl, amino, cyano, haloalkoxy, haloalkyl, halogen.

30

13. A compound according to claim 12 selected from the group consisting of

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-cyanobenzenesulfonamide;
4-cyano-N-((3R)-1-{3-[4-
(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;
5 N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;
4-bromo-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
fluorobenzenesulfonamide;
10 4-chloro-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;
N-(4-{[(3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl]amino}sulfonyl)phenyl)acetamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
15 methoxybenzenesulfonamide;
4-tert-butyl-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
methylbenzenesulfonamide;
20 N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
(trifluoromethyl)benzenesulfonamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2,5-
dimethoxybenzenesulfonamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-
25 methylbenzenesulfonamide;
3-chloro-N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
fluorobenzenesulfonamide;
N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
ethylbenzenesulfonamide;
30 N-((3R)-1-{3-[4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
isopropylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide;

2-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

5 3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

3,5-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

10 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

3-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide;

15 4-butoxy-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide;

20 3-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide;

25 2-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide;

30 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2,4-difluorobenzenesulfonamide;

5 N-(2-chloro-4-{[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl]amino}sulfonyl)phenyl)acetamide;

3,4-dichloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-

10 (cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

4-bromo-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

15 4-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methoxybenzenesulfonamide;

4-tert-butyl-N-((3R)-1-{3-[4-

20 (cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-(trifluoromethyl)benzenesulfonamide;

25 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2,5-dimethoxybenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-methylbenzenesulfonamide;

30 3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-fluorobenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-ethylbenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-isopropylbenzenesulfonamide;

5 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-fluorobenzenesulfonamide;

2-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

10 3-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-4-

15 (trifluoromethoxy)benzenesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-2,4-difluorobenzenesulfonamide;

N-((4-[(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl]amino)sulfonyl)-2-chlorophenyl)acetamide;

20 N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-3,4-dichlorobenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,5-bis(trifluoromethyl)benzenesulfonamide;

4-butoxy-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

25 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3,4-dimethoxybenzenesulfonamide;

3-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-methylbenzenesulfonamide;

30 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(trifluoromethoxy)benzenesulfonamide;

2-cyano-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-methylbenzenesulfonamide;

5 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-(trifluoromethyl)benzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-fluorobenzenesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-4-(trifluoromethoxy)benzenesulfonamide;

10 N-(2-chloro-4-{[(3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl]amino}sulfonyl}phenyl)acetamide;

3,4-dichloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

15 4-methoxy-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

4-isopropyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

4-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

20 3-cyano-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

3,4-dimethoxy-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)benzenesulfonamide;

25 N-(2-chloro-4-{[(3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl]amino}sulfonyl}phenyl)acetamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-fluoro-N-methylbenzenesulfonamide;

30 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-isopropyl-N-methylbenzenesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-(methylsulfonyl)benzenesulfonamide; and

5 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-(methylsulfonyl)benzenesulfonamide.

14. A compound according to claim 8 wherein

Z is CH₂;

10 R₃ is selected from the group consisting of hydrogen and alkyl;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

15 R₉ is hydrogen; and

R₁₀ is arylalkenyl wherein the aryl portion of said arylalkenyl is phenyl.

20 15. A compound according to claim 14 selected from the group consisting of

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-phenylethenesulfonamide; and

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-phenylethenesulfonamide.

25

16. A compound according to claim 8 wherein

Z is CH₂;

R₃ is selected from the group consisting of hydrogen and alkyl;

30 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-

morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

R₉ is hydrogen; and

5 R₁₀ is heterocycle wherein said heterocycle is selected from the group consisting of benzothienyl, imidazolyl, isoquinolinyl, pyridinyl, pyrrolyl, quinolinyl, thiazolyl, and thienyl, wherein said benzothienyl, imidazolyl, isoquinolinyl, pyridinyl, pyrrolyl, quinolinyl, thiazolyl, and thienyl are each optionally substituted with 1 or 2 substituents selected from the group consisting of alkoxy, alkyl, amino, haloalkyl, and halogen.

10

17. A compound according to claim 16 selected from the group consisting of

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-

15 thiophenesulfonamide;

5-chloro-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3-methyl-1-benzothiophene-2-sulfonamide;

N-(5-{[(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-

yl)oxy]propyl}pyrrolidinyl]amino]sulfonyl}-4-methyl-1,3-thiazol-2-yl)acetamide;

20 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-8-quinolinesulfonamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-7-isoquinolinesulfonamide;

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-isoquinolinesulfonamide;

25 N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-1-methyl-1H-imidazole-4-sulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-thiophenesulfonamide;

30 5-chloro-N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-3-methyl-1-benzothiophene-2-sulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-quinolinesulfonamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-8-isoquinolinesulfonamide; and

5 1-methyl-N-((3R)-1-{3-[4-(2-pyridinyl)phenoxy]propyl}pyrrolidinyl)-1H-imidazole-4-sulfonamide.

18. A compound according to claim 8 wherein

Z is CH₂;

10 R₃ is selected from the group consisting of hydrogen and alkyl;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-

15 azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

R₉ is hydrogen; and

R₁₀ is heterocycleheterocycle wherein said heterocycleheterocycle is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)-1H-pyrrol-2-yl.

20

19. A compound according to claim 18 that is N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-1-[3-chloro-5-(trifluoromethyl)-2-pyridinyl]-1H-pyrrole-2-sulfonamide.

25 20. A compound according to claim 8 wherein

Z is CH₂;

R₃ is selected from the group consisting of hydrogen and alkyl;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-

azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;


R₉ is hydrogen; and

R₁₀ is heterocycloxyaryl wherein said heterocycloxyaryl is (3-chloro-5-(trifluoromethyl)-2-pyridinyl)oxy)phenyl).

21. A compound according to claim 20 that is N-[(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl]-4-[(3-chloro-5-(trifluoromethyl)-2-pyridinyl)oxy]benzenesulfonamide.

10

22. A compound according to claim 1 of formula III



III,

or a pharmaceutically acceptable salt thereof wherein

15

R₁₁ is selected from the group consisting of hydrogen, alkenyl, alkenyloxy, alkoxy, alkyl, alkynyl, alkynyoxy, amino, aryl, arylalkenyl, arylalkyl, arylaryl, arylheterocycle, aryloxyaryl, cycloalkyl, cycloalkylalkyl, heterocycle, heterocyclealkyl, heterocyclearyl, heterocycleheterocycle, heterocycloxyalkyl, heterocycloxyaryl and heterocyclethioalkyl.

20

23. A compound according to claim 22 wherein

R₃ is selected from the group consisting of hydrogen, alkyl, alkylcarbonyl, aminocarbonyl, arylcarbonyl, and heterocyclecarbonyl;

R₈ is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle; and

25

R₁₁ is selected from the group consisting of alkenyloxy, alkoxy, alkyl, amino, aryl, cycloalkyl, heterocycle, heterocycleheterocycle and heterocyclethioalkyl.

24. A compound according to claim 22 wherein

Z is CH₂;

R₃ is selected from the group consisting of hydrogen, alkyl, alkylcarbonyl, and aminocarbonyl;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-5 cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

10 R₉ is hydrogen; and

R₁₁ is selected from the group consisting of alkenyloxy, alkoxy, alkyl, amino, and cycloalkyl.

25. A compound according to claim 24 selected from the group consisting of

15 N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}cyclopropanecarboxamide; tert-butyl (3R)-1-{2-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]ethyl}pyrrolidinylcarbamate; tert-butyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;

20 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylacetamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,3,3-trimethylbutanamide;

25 methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;

tert-pentyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl(methyl)carbamate;

30 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N',N'-trimethylurea;

N'-tert-butyl-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylurea;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)acetamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-3,3-dimethylbutanamide; 5
allyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate; methyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate; tert-pentyl (3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylcarbamate; 10
N'-(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N,N-dimethylurea; N-(tert-butyl)-N'-(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinylurea; 15
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(3,3-dimethylbutanoyl)-3,3-dimethylbutanamide; and
N'-(3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N'-(dimethylaminocarbonyl)-N,N-dimethylurea.

15

26. A compound according to claim 22 wherein

Z is CH_2 ;

R₃ is selected from the group consisting of hydrogen, alkyl, and arylcarbonyl wherein the aryl portion of said arylcarbonyl is phenyl optionally substituted with 1 substituent selected from the group consisting of cyano and halogen;

20 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

25 R₉ is hydrogen; and

R₁₁ is aryl wherein said aryl is phenyl optionally substituted with 1 substituent selected from the group consisting of cyano and halogen.

30

27. A compound according to claim 26 selected from the group consisting of

N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}-4-cyanobenzamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-
methylbenzamide;
4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-
5 methylbenzamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-fluoro-N-(4-
fluorobenzoyl)benzamide;
N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-
fluorobenzamide; and
10 4-cyano-N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-
yl)oxy]propyl}pyrrolidinyl)benzamide.

28. A compound according to claim 22 wherein
Z is CH₂;
15 R₃ is selected from the group consisting of hydrogen, alkyl, and heterocyclecarbonyl
wherein the heterocycle portion of said heterocyclecarbonyl is selected from the group
consisting of 4-morpholinyl and 1-pyrrolidinyl;
R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-
cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-
20 piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-
morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-
azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-
piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;
R₉ is hydrogen; and
25 R₁₁ is heterocycle wherein said heterocycle is selected from the group consisting of
furyl, 4-morpholinyl, pyridinyl, and pyrrolidinyl.
29. A compound according to claim 28 selected from the group consisting of
N-{(3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl}nicotinamide;
30 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-1-
pyrrolidinecarboxamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-4-morpholinecarboxamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methylnicotinamide;

5 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-furamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-1-pyrrolidinecarboxamide;

10 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-4-morpholinecarboxamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)nicotinamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-furamide;

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(1-pyrrolidinylcarbonyl)-1-pyrrolidinecarboxamide; and

15 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-(4-morpholinylcarbonyl)-4-morpholinecarboxamide.

30. A compound according to claim 22 wherein

Z is CH₂;

20 R₃ is selected from the group consisting of hydrogen and alkyl;

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;

25 R₉ is hydrogen; and

R₁₁ is heterocycleheterocycle wherein said heterocycleheterocycle is 2-(3-pyridinyl)-1,3-thiazol-4-yl.

30

31. A compound according to claim 30 selected from the group consisting of

N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide;

N-((3R)-1-{3-[4-(cyclopropylcarbonyl)phenoxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide;

5 N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-N-methyl-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide; and

N-((3R)-1-{3-[(4'-cyano[1,1'-biphenyl]-4-yl)oxy]propyl}pyrrolidinyl)-2-(3-pyridinyl)-1,3-thiazole-4-carboxamide.

10 32. A compound according to claim 22 wherein

Z is CH_2 ;

R_3 is selected from the group consisting of hydrogen and alkyl;

15 R_8 is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl;


R_9 is hydrogen; and

20 R_{11} is heterocyclethioalkyl wherein said heterocyclethioalkyl is [(4-methyl-2-pyrimidinyl)sulfanyl]methyl.

33. A compound according to claim 32 that is N-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-2-[(4-methyl-2-pyrimidinyl)sulfanyl]acetamide.

25

34. A compound according to claim 1 of formula IV



IV,

or a pharmaceutically acceptable salt thereof wherein

R₃ and R₄ are independently selected from the group consisting of hydrogen, alkenyl, alkyl, alkynyl, arylalkyl, cycloalkyl, cycloalkylalkyl, heterocycle and heterocyclealkyl.

5 35. A compound according to claim 34 wherein

R₃ and R₄ are independently selected from the group consisting of hydrogen and alkyl;

R₈ is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

10

36. A compound according to claim 34 wherein

Z is CH₂;

R₃ and R₄ are independently selected from the group consisting of hydrogen and alkyl;

15

R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and R₉ is hydrogen.

20 37. A compound according to claim 36 selected from the group consisting of

1-(4-{3-[(3R)-3-aminopyrrolidinyl]propoxy}phenyl)-1-propanone;

25

4'-{3-[(3R)-3-aminopyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;


4'-{3-[(3R)-3-(dimethylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile;

and

4'-{3-[(3R)-3-(methylamino)pyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile.

30

38. A compound according to claim 1 of formula V



V,


or a pharmaceutically acceptable salt thereof.

5 39. A compound according to claim 38 wherein
 R₈ is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

10 40. A compound according to claim 38 wherein
 Z is CH₂;
 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and
 R₉ is hydrogen.

15 41. A compound according to claim 40 selected from the group consisting of
 cyclopropyl{4-[3-(3-hydroxy-1-pyrrolidinyl)propoxy]phenyl}methanone; cyclopropyl(4-{3-[(3R)-3-hydroxypyrrolidinyl]propoxy}phenyl)methanone; 4'-{3-[(3R)-3-hydroxypyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile; 4'-{3-[(3S)-3-hydroxypyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile; 4'-[3-(3-hydroxy-3-methyl-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile; 25 4'-[3-(3-hydroxy-3-isopropyl-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile; and 4'-{3-[(3R)-3-hydroxy-3-methylpyrrolidinyl]propoxy}[1,1'-biphenyl]-4-carbonitrile.

42. A compound according to claim 1 of formula VI



VI,

or a pharmaceutically acceptable salt thereof.

5

43. A compound according to claim 42 wherein

R8 is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

10

44. A compound according to claim 42 wherein

Z is CH₂;

R8 is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and


15

R9 is hydrogen.

20

45. A compound according to claim 44 that is 4'-[3-(3-oxo-1-pyrrolidinyl)propoxy][1,1'-biphenyl]-4-carbonitrile.

46. A compound according to claim 1 of formula VII



VII,

or a pharmaceutically acceptable salt thereof.

47. A compound according to claim 46 wherein

5 R₈ is selected from the group consisting of alkylcarbonyl, cycloalkylcarbonyl, aryl, heterocycle, heterocyclecarbonylaryl and heterocyclecarbonylheterocycle.

48. A compound according to claim 46 wherein

Z is CH₂;10 R₈ is selected from the group consisting of acetyl, propionyl, cyclopropylcarbonyl, 4-cyanophenyl, 2-pyridinyl, 4-(1-pyrrolidinylcarbonyl)phenyl, 4-(1-pyrrolidinylcarbonyl)-1-piperazinyl, 4-(1-piperidinylcarbonyl)phenyl, 4-(1-piperidinylcarbonyl)-1-piperazinyl, 4-(4-morpholinylcarbonyl)phenyl, 4-(4-morpholinylcarbonyl)-1-piperazinyl, 4-(1-azetidinylcarbonyl)phenyl, 4-(1-azetidinylcarbonyl)-1-piperazinyl, 4-(1-piperazinylcarbonyl)phenyl, 4-(1-piperazinylcarbonyl)-1-piperazinyl; and15 R₉ is hydrogen.

49. A compound according to claim 46 selected from the group consisting of

20 (5R)-3-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-methyl-2,4-imidazolidinedione;

(5S)-3-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-methyl-2,4-imidazolidinedione; and

(5R)-3-((3R)-1-[3-(4-acetylphenoxy)propyl]pyrrolidinyl)-5-isopropyl-2,4-imidazolidinedione.

25

50. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 1 in combination with a pharmaceutically acceptable carrier.

30

51. A method for modulating the effects of the histamine-3 receptor in a mammal comprising administering a therapeutically effective amount of a compound of claim 1.

52. A method of treating a disorder wherein the disorder is ameliorated by modulating the histamine-3 receptor in a mammal in need of such treatment comprising administering a therapeutically effective amount of a compound of claim 1.

5 53. The method of claim 52 wherein the disorder is selected from the group consisting of acute myocardial infarction, asthma, cutaneous carcinoma, depression, inflammation, medullary thyroid carcinoma, melanoma, Meniere's disease, migraine, motion sickness, obesity, pain, Parkinson's disease, schizophrenia, seizures, and septic shock.

10 54. The method of claim 52 wherein the disorder is Alzheimer's disease.

55. The method of claim 52 wherein the disorder is attention-deficit hyperactivity disorder.

15 56. The method of claim 52 wherein the disorder is epilepsy.

57. The method of claim 52 wherein the disorder is narcolepsy.

INTERNATIONAL SEARCH REPORT

In International Application No
PCT/US 01/21929

A. CLASSIFICATION OF SUBJECT MATTER					
IPC 7	C07D207/12	C07D207/14	C07D207/16	C07D401/12	C07D401/14
	C07D403/12	C07D405/12	C07D409/12	C07D417/12	C07D403/04
	C07D417/14	A61K31/4025	A61P9/10	A61P25/28	A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 49706 A (ALTMANN EVA ;CIBA GEIGY AG (CH); MISSBACH MARTIN (CH); WIDLER LEO) 31 December 1997 (1997-12-31) claims 6,7; examples 91,102	1,38,50
X	EP 0 738 725 A (LILLY CO ELI) 23 October 1996 (1996-10-23) claims 1,2,4; examples 8,13,64,65	1,38,50
A	EP 0 978 512 A (BIOPROJET SOC CIV) 9 February 2000 (2000-02-09) cited in the application the whole document	1-57

 Further documents are listed in the continuation of box C.

Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the International filing date
- *L* document which may throw doubts on priority, claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the International filing date but later than the priority date claimed

T later document published after the International filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

Y document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Z document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

Date of mailing of the International search report

22 November 2001

03/12/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

Int'l. Application No

PCT/US 01/21929

Patent document cited in search report	Publication date		Patent family member(s)	Publication date
WO 9749706	A 31-12-1997	AU WO	3176297 A 9749706 A1	14-01-1998 31-12-1997
EP 0738725	A 23-10-1996	AU CA EP JP WO	5554996 A 2215902 A1 0738725 A2 11504013 T 9632937 A1	07-11-1996 24-10-1996 23-10-1996 06-04-1999 24-10-1996
EP 0978512	A 09-02-2000	EP AU WO EP EP	0978512 A1 5511999 A 0006254 A2 0982300 A2 1100503 A2	09-02-2000 21-02-2000 10-02-2000 01-03-2000 23-05-2001